

CALDERON PROJECTOR FOR THE HESSIAN OF THE PERTURBED CHERN-SIMONS FUNCTION ON A 3-MANIFOLD WITH BOUNDARY

BENJAMIN HIMPEL, PAUL KIRK, AND MATTHIAS LESCH

ABSTRACT. The existence and continuity for the Calderón projector of the perturbed odd signature operator on a 3-manifold is established. As an application we give a new proof of a result of Taubes relating the mod 2 spectral flow of a family of operators on a homology 3-sphere with the difference in local intersection numbers of the character varieties coming from a Heegard decomposition.

1. INTRODUCTION

The purpose of this article is to establish the existence and continuity of the Calderón projectors for families of perturbed signature operators on a 3-manifold with boundary. We explain briefly the motivation.

The Morse theory of the Chern-Simons function on the space of $SU(n)$ connections over a 3-manifold leads to the construction of topological invariants, notably Taubes' construction of Casson's invariant [29], Floer's instanton homology [17], and more recently $SU(3)$ and $SU(n)$ extensions of Casson's invariant [3, 4, 11, 19]. These invariants are defined using the Morse index of the Hessian of the Chern-Simons function at a critical point. To make this rigorous Taubes showed (in the $SU(2)$ setting) how to understand this ill-defined Morse index in terms of the spectral flow of the odd signature operator coupled to a path of connections. Most importantly for the present article, he also showed how to construct perturbations of the Chern-Simons function in order to make it suitably non-degenerate.

Taubes' ideas form the blueprint for generalizations of Casson's invariant. In particular the definition of the generalized Casson invariants involve the spectral flow of 1-parameter families of perturbed odd signature operators. These perturbed odd signature operators (which we describe in detail below) are compact perturbations of Dirac operators, but they are not differential operators.

With the construction of a Morse-theoretic topological invariant established, the problem arises of computing the invariant to discover what information it contains. Casson's surgery formula and Floer's exact triangle are examples of computational formulas which make it possible to compute their invariants. In general one wishes to have surgery formulas for the generalized Casson invariants. This leads to the problem of computing the spectral flow of the Hessian of the Chern-Simons function between critical points in terms of cut-and-paste data for the underlying 3-manifold.

There is a good cut-and-paste theory for the spectral flow of Dirac operators which make it possible to carry out calculations. Nicolaescu's article [25] gives a splitting formula for the spectral flow of a family of Dirac operators. This formula equates the spectral flow to an infinite-dimensional Maslov index corresponding to the associated paths of Calderón projectors coming from two parts in a decomposition of the manifold. (Roughly speaking, the Calderón projector for a Dirac operator D is the projection in $L^2(\partial X)$ onto the subspace of boundary values of solutions to the equation $D\phi = 0$.) Nicolaescu's result, combined with some techniques introduced in [12], sufficed to compute $SU(3)$ Casson invariants for certain Seifert fibered homology 3-spheres in [5, 11], where one could avoid using perturbations (at least near the reducible critical points). However, the presence of

Date: Feb. 18, 2003.

The second named author gratefully acknowledges the support of the National Science Foundation under grant no. DMS-9971020.

singularities in the flat moduli space complicates calculations for more general homology spheres, requiring the use of perturbations. To apply the powerful methods of [25] one needs an extension of the results concerning the existence and continuity of the Calderón projectors for families of operators more general than Dirac operators; in fact general enough to include the perturbed odd signature operator. This is the task accomplished in this article. Our main result is the following, which combines Theorems 3.16 and 3.8.

Theorem. *Consider the class $\mathcal{D}$ of Dirac operators on $E \rightarrow X$ for a manifold with boundary X which are in cylindrical form near ∂X , and the class $\mathcal{V}$ of auxiliary operators satisfying the pseudolocality property (3.6). Then for $D \in \mathcal{D}$ and $V \in \mathcal{V}$ the Calderón projector of $D + V$ is well defined.*

We say that a family $D_t \in \mathcal{D}$ varies continuously if all coefficients of D_t (in any local chart) vary continuously. Moreover, we equip $\mathcal{V}$ with the usual norm topology of $\mathcal{B}(L^2(X, E))$. Then for each $n \in \mathbb{Z}_+$ the map

$$\{(D, V) \in \mathcal{D} \times \mathcal{V} \mid \dim Z_0(D + V) = n\} \longrightarrow \mathcal{B}(L^2(\Sigma, E)), \quad (D, V) \mapsto C(D + V)$$

taking a pair (D, V) to the Calderón projector C for the operator $D + V$ is continuous.

In the statement, $Z_0(D + V)$ is a vector space isomorphic to $K \oplus K$, where K denotes the space of sections ϕ which vanish on a collar of the boundary and which satisfy $(D + V)\phi = 0$. For Dirac operators the unique continuation property implies that $Z_0 = 0$. A novel feature of our approach is that we do not require the unique continuation property to hold.

We are chiefly concerned with the application of this result to the perturbed odd signature operator arising from the Morse theory of the Chern-Simons function, as this will be used crucially in a forthcoming article [6]. (In fact the completion of that article was held up by the need for the main result of the present article.) Unfortunately there is no suitably precise exposition of the various constructions which arise in this context and so we include the details necessary to set up the problem precisely and also for the convenience of the reader. One of our purposes is to explicitly provide justification for comments like “in the non-generic case one must perturb, but the arguments carry through,” in the context of cut-and-paste techniques for spectral flow.

It is our hope that in addition to providing the proof of the theorem stated above this article complements the exposition [18, 19]. Thus in Section 2 we explain the approach used in the Morse-theoretic study of the Chern-Simons function, with an emphasis on how the perturbations are constructed and on the effect of perturbing on the Hessian of the Chern-Simons function. We also include proofs of some (known) results concerning the pseudolocality of the perturbation (Proposition 2.2) and an estimate concerning the size of the perturbation (Lemma 2.3) that we require to apply the result stated above to the context of the Hessian of the perturbed Chern-Simons function.

Theorem 3.19 is our desired application. In this statement, $D_{A,f}$ denotes the perturbed twisted odd signature operator coupled to a connection A and a perturbation f . It is essentially the Hessian of the perturbed Chern-Simons function $cs + f : \mathcal{A} \rightarrow \mathbb{R}$.

Theorem 3.19. *Let X be a compact 3-manifold with boundary and $\mathcal{A}$ the space of connections in a principal $SU(n)$ bundle over X . Then there exists a neighborhood $U \subset \mathcal{A}$ of the space of flat connections, and an $\epsilon > 0$ so that if $B_{\mathcal{P}}(0, \epsilon)$ denotes the ϵ ball around 0 in the space of perturbations $\mathcal{P}$, the map $U \times B_{\mathcal{P}}(0, \epsilon) \rightarrow \mathcal{B}(L^2(\Sigma, E))$ sending a pair (A, f) to the Calderón projector $P_{A,f}$ of $D_{A,f}$ is continuous.*

We finish the article by giving a new proof in Section 4, using our methods, of the main step in Taubes’ argument that his invariant equals Casson’s invariant.

Theorem (Taubes). *The mod 2 Spectral flow of the perturbed odd signature operator between two generic perturbed flat connections on a homology 3-sphere equals zero if and only if the corresponding*

local intersection numbers of the $SU(2)$ character varieties of the Heegard handlebodies in the $SU(2)$ character variety of the Heegard surface are equal.

The strategy of our proof is to use an adiabatic stretching of a collar of the Heegard surface and symplectic reduction to reduce the infinite-dimensional Maslov index (which equals the sign Taubes assigns to a critical point) to a finite-dimensional Maslov index in the character variety of a Heegard surface (which determines Casson's sign). Having established in Theorem 3.19 that the Calderón projectors exist and vary nicely, the first step in the argument uses this and Nicolaescu's theorem [25] to identify spectral flow with the infinite-dimensional Maslov index of the Calderón projectors. The rest of the argument is a study of the effect of stretching and a corresponding symplectic reduction, as analyzed in [12].

The authors thank H. Boden, G. Daskalopoulos, C. Herald, T. Mrowka, and K. Wojciechowski for helpful discussions.

2. BASIC SET-UP

2.1. Connections on a 3-manifold. We begin by recalling the spaces and operators of interest.

Let X be a compact 3-manifold. Fix a principal $SU(n)$ bundle P over X . We associate two vector bundles to P : the $\mathbb{C}^n$ vector bundle $E \rightarrow X$ associated to the standard representation $SU(n) \rightarrow GL(\mathbb{C}^n)$, and the Lie algebra bundle $\text{ad } P \rightarrow X$ associated to the adjoint representation $SU(n) \rightarrow GL(\mathfrak{su}(n))$ of $SU(n)$ on its Lie algebra $\mathfrak{su}(n)$. Notice that the bundle P is trivializable since $\pi_1(SU(n))$ and $\pi_2(SU(n))$ are trivial.

We let $\mathcal{A}$ denote the space of smooth connections on P . A connection $A \in \mathcal{A}$ defines a covariant derivative

$$d_A : C^\infty(X; E) \rightarrow \Omega^1(X; E).$$

It is convenient to view $\text{ad } P$ as a subbundle of $\text{End}(E)$. Then the covariant derivative on $\text{ad } P$ associated to A (which we also denote by $d_A : C^\infty(X; \text{ad } P) \rightarrow \Omega^1(X; \text{ad } P)$) satisfies

$$(d_A(\phi))(e) = d_A(\phi(e)) - \phi(d_A(e)) \quad \text{for } \phi \in C^\infty(X; \text{ad } P), \quad e \in C^\infty(X; E).$$

The covariant derivatives extend to p -forms $d_A : \Omega^p(X; E) \rightarrow \Omega^{p+1}(X; E)$ and $d_A : \Omega^p(X; \text{ad } P) \rightarrow \Omega^{p+1}(X; \text{ad } P)$ by requiring the graded Leibnitz rule to hold. The operator $d_A^2 : \Omega^p(X; E) \rightarrow \Omega^{p+2}(X; E)$ has order zero and is given by multiplication by the curvature form $F(A) \in \Omega^2(X; \text{ad } P)$.

The space $\mathcal{A}$ is an affine space modeled on the space of Lie algebra valued 1-forms $\Omega^1(X; \text{ad } P)$: Fixing a connection A_0 yields an identification

$$(2.1) \quad \mathcal{A} \xrightarrow{\cong} \Omega^1(X; \text{ad } P), \quad A \mapsto A - A_0.$$

It is sometimes convenient to fix a trivialization $P \cong X \times SU(n)$; then we can take A_0 to be the trivial (product) connection which we denote by Θ . The space $\mathcal{A}$ is acted on by the group of (smooth) gauge transformations $\mathcal{G} = C^\infty(X; P \times_{\text{Ad}} SU(n))$ in the usual way: $d_{gA}(e) = g(d_A(g^{-1}(e)))$.

We fix a Riemannian metric on X . If the boundary of X is non-empty we will always assume that a collar of the boundary ∂X in X is isometric to $[0, 1) \times \partial X$.

The L^2 inner product on $\Omega^i(X; \text{ad } P)$ is then defined in terms of the Hodge $*$ operator on X by the formula

$$(2.2) \quad \langle b, c \rangle_{L^2} = - \int_X \text{tr}(b \wedge *c).$$

For analytical reasons one must at times complete the spaces $\mathcal{A}$ and $\mathcal{G}$ in appropriate Sobolev norms. Thus we fix some $s \geq 1$ and take $\mathcal{A}_s$ to be the L_s^2 completion of $\mathcal{A} \cong \Omega^1(X; \text{ad } P)$, and $\mathcal{G}_{s+1}$ to be the closure of $\mathcal{G}$ in the L_{s+1}^2 completion of $C^\infty(\text{End}(E))$. Here L_s^2 denotes the completion in the Sobolev norm corresponding to sections with s derivatives in L^2 , extended in the standard way to all $s \in \mathbb{R}$. Then $\mathcal{G}_{s+1}$ acts smoothly on $\mathcal{A}_s$ (see [20]). The restriction $s \geq 1$ guarantees that the gauge transformations are continuous.

2.2. The Chern-Simons function. The Chern-Simons function $cs : \mathcal{A} \rightarrow \mathbb{R}$ is defined in terms of a trivialization $P \cong X \times SU(n)$ as follows. To a connection $A \in \mathcal{A}$ one can assign the $su(n)$ -valued 1-form

$$a := A - \Theta \in \Omega^1(X; \text{ad } P) \cong \Omega^1(X) \otimes su(n).$$

One then defines

$$cs(A) = \frac{1}{8\pi^2} \int_X \text{tr}(da \wedge a + \frac{2}{3}a \wedge a \wedge a).$$

In this expression (as in Equation (2.2)) the wedge product means the wedging of the differential forms and matrix multiplication of the $su(n)$ coefficients. Note that the definition (and value) of the Chern-Simons function depends on the choice of trivialization of P . If X is *closed*, then the reduction of $cs(A)$ modulo $\mathbb{Z}$ is independent of the choice of trivialization, and is gauge invariant. In fact, $cs(g \cdot A) = cs(A) + \text{degree}(g)$, where the degree is defined using the fact that $H^3(SU(n); \mathbb{Z}) = \mathbb{Z}$. Thus when X is closed, one considers the Chern-Simons function as a circle valued function on the quotient $cs : \mathcal{A}/\mathcal{G} \rightarrow \mathbb{R}/\mathbb{Z} = S^1$.

In [29], Taubes gave an interpretation of Casson's invariant of homology 3-spheres by studying the Morse theory of the Chern-Simons function. This was generalized in much subsequent work, including the work of Floer [17] and Herald-Boden [3]. Thus we look next at the differential topology of $\mathcal{A}$, $\mathcal{G}$, and cs .

The tangent space of $\mathcal{A}$ at a connection A is canonically identified with $\Omega^1(X; \text{ad } P)$. It is not hard to identify the tangent space of $\mathcal{G}$ at the identity with $\Omega^0(X; \text{ad } P)$. Fixing a connection $A \in \mathcal{A}$ yields the *action map*

$$g_A : \mathcal{G} \rightarrow \mathcal{A}, \quad g \mapsto g \cdot A$$

The differential of this map is just the map $d_A : \Omega^0(X; \text{ad } P) \rightarrow \Omega^1(X; \text{ad } P)$. Precisely, the diagram

$$(2.3) \quad \begin{array}{ccc} T_1 \mathcal{G} & \xrightarrow{dg_A} & T_A \mathcal{A} \\ \cong \downarrow & & \cong \downarrow \\ \Omega^0(X; \text{ad } P) & \xrightarrow{d_A} & \Omega^1(X; \text{ad } P) \end{array}$$

commutes. Thus the tangent space to the quotient $\mathcal{A}/\mathcal{G}$ at $[A]$, $T_{[A]} \mathcal{A}/\mathcal{G}$ can be identified with the cokernel of d_A (at least at smooth points of $\mathcal{A}/\mathcal{G}$: making this precise requires looking at Sobolev completions and dealing with quotient singularities but we will not emphasize this point here).

2.3. The Hessian of the Chern-Simons function. *In this section we assume that X is a closed 3-manifold.* This is because the differential operators which arise from the Morse theory of the Chern-Simons function have a simple description on a closed manifold which is adequate for our purposes, even though we are ultimately interested in cutting and pasting questions. This section could be written in the context of manifolds with boundary, but in that case the proper way to study the Chern-Simons function is as a section of a determinant bundle as in [27] and [18]. (The material of this subsection is well known. Standard references include [29, 15, 20]. We include it to set up notation and to prepare the reader for the section on perturbations.)

The differential $dcs_A : T_A \mathcal{A} \rightarrow \mathbb{R}$ of the Chern-Simons function is computed (see [29]) for X *closed* as:

$$dcs_A(b) = \frac{1}{4\pi^2} \int_X \text{tr}(F(A) \wedge b),$$

with $F(A)$ the curvature 2-form of A . (Recall that in a trivialization, $F(A) = da + a \wedge a$.) From this one concludes that the critical points of $cs : \mathcal{A} \rightarrow \mathbb{R}$ are those A such that $F(A) = 0$, i.e. the *flat connections*.

The Hessian of $cs : \mathcal{A} \rightarrow \mathbb{R}$, $\text{Hess}_A(cs) : T_A \mathcal{A} \times T_A \mathcal{A} \rightarrow \mathbb{R}$ is defined for any connection $A \in \mathcal{A}$ using the affine structure of $\mathcal{A}$. It can be computed explicitly when X is closed ([29]):

$$\text{Hess}(cs)_A(b, c) = -\frac{1}{4\pi^2} \langle *d_A b, c \rangle.$$

The L^2 inner product can be viewed as a Riemannian metric on the infinite-dimensional manifold $\mathcal{A}$ (or $\mathcal{A}_s$). Identifying vectors and covectors via this metric allows us to define the gradient vector field of cs by the rule $\langle \text{grad}(cs)_A, b \rangle_{L^2} = dcs_A(b)$. Similarly the linearization $H_A : T_A \mathcal{A} \rightarrow T_A \mathcal{A}$ of $\text{grad}(cs)$ is obtained from the Hessian by the rule $\langle H_A(b), c \rangle_{L^2} = \text{Hess}(cs)_A(b, c)$. From the above formulas one concludes that

$$\text{grad}(cs)_A = -\frac{1}{4\pi^2} * F(A)$$

and

$$H_A(b) = -\frac{1}{4\pi^2} * d_A(b).$$

The self-adjoint operator H_A is not elliptic because of the gauge invariance of cs . For example, at a flat connection A , $d_A d_A = F(A) = 0$, and so $H_A(d_A(b)) = 0$ for all $b \in \Omega^0(X; \text{ad } P)$. In other words the kernel of H_A contains the infinite-dimensional subspace of tangent vectors to the $\mathcal{G}$ orbit through A . This causes problems if one wishes to consider the Morse theory of cs .

As explained in [29] (and is typical in gauge theory), the correct way to resolve this lack of ellipticity is to consider $\text{grad}(cs)$ as a vector field on the quotient $\mathcal{A}/\mathcal{G}$, and thus H_A as a family of elliptic self-adjoint operators on $T_{[A]} \mathcal{A}/\mathcal{G}$. Using the slice theorem and Diagram (2.3) one identifies $T_{[A]} \mathcal{A}/\mathcal{G}$ (one has to take care at the singular points) as the orthogonal complement of the image of $d_A : \Omega^0(X; \text{ad } P) \rightarrow \Omega^1(X; \text{ad } P)$, in other words as the kernel of the adjoint $d_A^* : \Omega^1(X; \text{ad } P) \rightarrow \Omega^0(X; \text{ad } P)$. The Bianchi identity implies that $d_A^*(\text{grad}(cs)(A)) = -\frac{1}{4\pi^2} d_A^*(F(A)) = 0$. If A is a critical point of cs (i.e. a flat connection), then

$$d_A^*(H_A(b)) = \frac{1}{4\pi^2} * d_A * d_A(b) = \frac{1}{4\pi^2} * d_A d_A(b) = \frac{1}{4\pi^2} * F(A)(b) = 0.$$

But $H_A(b)$ is not necessarily in the kernel of d_A^* when A is not a critical point. This motivates setting

$$\tilde{H}_A : \ker d_A^* \rightarrow \ker d_A^*, \quad \tilde{H}_A(b) = -\frac{1}{4\pi^2} \text{Proj}_{\ker d_A^*} (*d_A(b))$$

where Proj denotes the L^2 orthogonal projection. The operator $\tilde{H}_A$ agrees with the restriction of H_A to $\ker d_A^*$ if the connection A is flat.

Then $\tilde{H}_A$ is Fredholm (i.e. its extension to $L_s^2(\Omega^1(X; \text{ad } P)) \cap \ker d_A^* \rightarrow L_{s-1}^2(\Omega^1(X; \text{ad } P)) \cap \ker d_A^*$ is Fredholm). Moreover, it is self-adjoint with compact resolvent with respect to the L^2 inner product restricted to $\ker d_A^*$. The spaces $\ker d_A^*$ form a smooth vector bundle over each orbit-type stratum of $\mathcal{A}/\mathcal{G}$. In particular, Taubes' idea was to use the spectral flow of a family of operators $\tilde{H}_{A_t}$ as a substitute for the difference in Morse index of critical points A_0 and A_1 of the Chern-Simons function, and thereby to define an Euler characteristic for $\mathcal{A}/\mathcal{G}$.

(The *spectral flow* $\text{SF}(H_t)$ of a 1-parameter family of self-adjoint operators H_t is the integer defined to be the number of eigenvalues that cross zero, counted with sign. Some conventions need to be set if the operators in question have kernel at the endpoints of the path, and one needs to establish that the concept is well defined for appropriately continuous paths of unbounded operators. A careful construction appropriate for our purposes can be found in [8]. The basic properties and conventions are explained in [22].)

Taubes observed that the spectral flow of a family $\tilde{H}_{A_t}$ equals the spectral flow of a slightly better behaved family of operators. Consider, for any connection A , the twisted de Rham sequence

$$(2.4) \quad 0 \rightarrow \Omega^0(X; \text{ad } P) \xrightarrow{d_A} \Omega^1(X; \text{ad } P) \xrightarrow{d_A} \Omega^2(X; \text{ad } P) \xrightarrow{d_A} \Omega^3(X; \text{ad } P) \rightarrow 0.$$

The sequence (2.4) is an elliptic complex when A is flat. It is not a complex at a non-flat connection, but can be made into a complex if one substitutes $d'_A := -4\pi^2 * \tilde{H}_A \circ \text{Proj}_{\ker d_A^*}$ for $d_A : \Omega^1(X; \text{ad } P) \rightarrow \Omega^2(X; \text{ad } P)$. When A is flat, $d'_A = d_A$.

Whether or not A is flat, the *odd signature operator* (obtained essentially by folding up the sequence (2.4))

$$(2.5) \quad \begin{aligned} D_A &: \Omega^0(X; \text{ad } P) \oplus \Omega^1(X; \text{ad } P) \rightarrow \Omega^0(X; \text{ad } P) \oplus \Omega^1(X; \text{ad } P) \\ D_A(b, c) &= (d_A^* c, *d_A c + d_A b) \end{aligned}$$

is a self-adjoint Dirac operator in the sense of e.g. [10] for any connection A .

One also has a self-adjoint operator $D'_A(b, c) = (d_A^* c, *d'_A c + d_A b)$. For A flat, $D_A = D'_A$. For a general irreducible connection A , $D_A - D'_A$ is a relatively compact operator. In fact, letting $\Delta_A^{-1} : \Omega^0(X; \text{ad } P) \rightarrow \Omega^0(X; \text{ad } P)$ denote the inverse of $d_A^* d_A$ (which is invertible for A irreducible) and observing that $\text{Proj}_{\ker d_A^*} x = x - d_A \Delta_A^{-1} d_A^* x$, we see that

$$(D_A - D'_A)(b, c) = (*F(A)\Delta_A^{-1}d_A^* - d_A\Delta_A^{-1}*F(A) + d_A\Delta_A^{-1}*F(A)d_A\Delta_A^{-1}d_A^*)(c).$$

(see [29] for details).

By decomposing $\Omega^0(X; \text{ad } P) \oplus \Omega^1(X; \text{ad } P)$ as $\Omega^0(X; \text{ad } P) \oplus \text{Image } d_A \oplus \ker d_A^*$, one sees that the spectrum of D'_A is the disjoint union of the spectrum of $\tilde{H}_A$ and a symmetric (with respect to $\lambda \rightarrow -\lambda$) spectrum. A simple argument then shows that given a path A_t of (say irreducible) connections joining two critical points of cs , the spectral flow of the family $\tilde{H}_{A_t}$ (which plays the role of the difference in Morse indices of cs at A_0 and A_1) equals the spectral flow of the family D_{A_t} . What is gained is that the spectral flow of D_{A_t} makes sense whether or not A_t is a path of irreducible connections, i.e. even when the subspaces $\ker d_A^* \subset \Omega^1(X; \text{ad } P)$ do not vary continuously.

2.4. Perturbations. We come now to the construction of perturbations and their effect on the Hessian of the Chern-Simons function. The critical point set of cs may be complicated and it is often necessary to perturb the Chern-Simons function to make it as nice as possible.

Making precise what “nice” means can be quite difficult in the case of $SU(n)$ for $n > 2$, or when the homology of X is complicated. In the case of $SU(2)$ and X a homology sphere Taubes replaced cs by a function $cs + h$ whose restriction to the top stratum of $\mathcal{A}/\mathcal{G}$ (i.e. the irreducible connections; see Section 4) is Morse in a suitable sense, and then used this to define a topological invariant in the spirit of the Poincaré-Hopf theorem, which he showed equals Casson’s invariant. In [3] a similar analysis is carried out for $SU(3)$ and X a homology sphere leading to a definition of an $SU(3)$ Casson invariant; in that case in addition to being Morse on the top stratum of $\mathcal{A}/\mathcal{G}$, $cs + h$ must be suitably non-degenerate in higher codimensional strata. In more recent work [19] Herald shows how with a suitable choice of perturbation h one can define versions of Casson’s invariant for rational homology spheres and $SU(n)$, $n = 2, 3, 4, 5$.

There are slightly different approaches to constructing a suitable family of perturbations h , but they are similar and our method will apply to each approach. We mention that the basic requirements that h should satisfy are:

- (1) The perturbation h should be a smooth function on the Hilbert manifold $\mathcal{A}_1/\mathcal{G}_2$.
- (2) The operator defined by the Hessian of $cs + h$ at A should be a compact perturbation of $\tilde{H}_A$.
- (3) The family of admissible h should be large enough so that one can prove various general position results about $cs + h$ on the strata and their normal bundles of $\mathcal{A}/\mathcal{G}$.

The following types of perturbations, due to Taubes and Floer, accomplish these goals, as explained in [19].

First, fix a collection of smooth embeddings $\gamma_i : D^2 \times S^1 \rightarrow X$, $i = 1, \dots, N$. Taubes [29] requires that the embeddings have disjoint images, whereas Floer [17] and Herald [19] assume that $\gamma_i(x, 1) = \gamma_1(x, 1)$ for all i and $x \in D^2$ and that the derivatives of the γ_i in the S^1 direction at points in $D^1 \times \{1\}$ are independent of i . (In fact, nothing is lost by assuming that there is an interval I around $1 \in S^1$ so that $\gamma_i(x, u)$ is independent of i for $u \in I$.) The results of this article work equally well in each setting. The difference between them is that it is easier to show that the third condition above holds for perturbations of the type Floer considers since the base point issues are easier to

control. We will assume, to keep the notation under control, that we are using the approach of Floer, i.e. the common basepoint approach. We do not address the third condition in this article. In the language of [18, 19, 3] the third condition holds if the set of perturbations is *abundant*.

Next, let $\mathcal{P}$ denote the set of C^r functions (for some fixed large r), $f : SU(n)^N \rightarrow \mathbb{R}$, invariant under the conjugation action of $SU(n)$, i.e. $f(rg_1r^{-1}, \dots, rg_Nr^{-1}) = f(g_1, \dots, g_N)$.

Now let $P|_{D^2}$ denote the restriction of the principal bundle P to the disc $D^2 = \gamma_i(D^2 \times \{1\})$. Given a smooth connection $A \in \mathcal{A}$, consider the map $\Phi_A : P|_{D^2} \rightarrow SU(n)^N$ defined by the rule $\Phi_A(p) = (g_1, \dots, g_N)$ where $pg_i \in P_x$ is the endpoint of the parallel lift of the loop $\gamma_i(x, u)$, $u \in [0, 2\pi]$ with respect to A starting at p (where $x \in D^2$ is the image of p via the bundle projection). Then $\Phi_A(pr) = (r^{-1}g_1r, \dots, r^{-1}g_Nr)$ for $r \in SU(n)$. It follows that if $f \in \mathcal{P}$, $f \circ \Phi_A$ descends to a well-defined function on D^2 .

If the principal bundle is trivialized, $P \cong X \times SU(n)$, then one can view parallel lifting as the holonomy, a function from loops to $SU(n)$, and in our context this allows us to define

$$\text{Hol} : \mathcal{A} \times D^2 \rightarrow SU(n)^N, \quad (A, x) \mapsto (\text{Hol}_{\gamma_1(x, -)}(A), \dots, \text{Hol}_{\gamma_N(x, -)}(A)).$$

Thus we denote the function $f \circ \Phi_A$ by $x \mapsto f(\text{Hol}(A, x))$. Notice that $f(\text{Hol}(A, x))$ is independent of the trivialization, and is unchanged if we change A by a gauge transformation.

Fix a smooth positive cut-off function η on D^2 which vanishes near the boundary. The space of *admissible perturbations* is defined to be the space of functions of the form

$$(2.6) \quad h_f : \mathcal{A} \rightarrow \mathbb{R}, \quad A \mapsto \int_{D^2} f(\text{Hol}(A, x)) \eta(x) d^2x.$$

Thus we view the admissible perturbation functions as being parameterized by the (Fréchet) space $\mathcal{P}$.

Taubes and Floer show that h_f extends to a smooth function on $\mathcal{A}_2$. Note that $d(cs + h_f)_A = dcs_A + d(h_f)_A$ and $\text{Hess}(cs + h_f)_A = \text{Hess}(cs)_A + \text{Hess}(h_f)_A$. Thus one can define

$$Q_{A,f} : T_A \mathcal{A} \cong \Omega^1(X; \text{ad } P) \rightarrow T_A \mathcal{A}$$

by the rule

$$\langle Q_{A,f}(b), c \rangle = \text{Hess}(h_f)_A(b, c) = \left. \frac{\partial^2}{\partial t \partial s} \right|_{s=t=0} h_f(A + sb + tc).$$

Then setting $d_{A,f} := d_A - 4\pi^2 * Q_{A,f}$ we see that $-\frac{1}{4\pi^2} * d_{A,f}$ is the operator defined by the Hessian of $cs + h_f$ at A .

A connection A is called *f-perturbed flat* if it is a critical point of $cs + h_f$ or, equivalently, if $*F(A) = 4\pi^2 \text{grad}(h_f)_A$. As in Section 2.3, we form the sequence

$$(2.7) \quad 0 \rightarrow \Omega^0(X; \text{ad } P) \xrightarrow{d_A} \Omega^1(X; \text{ad } P) \xrightarrow{d_{A,f}} \Omega^2(X; \text{ad } P) \xrightarrow{d_A} \Omega^3(X; \text{ad } P) \rightarrow 0.$$

The sequence (2.7) is a complex if A is *f-perturbed flat*, and for any connection A and perturbation f the operator obtained from (2.7) (compare to (2.5)),

$$(2.8) \quad \begin{aligned} D_{A,f} : \Omega^0(X; \text{ad } P) \oplus \Omega^1(X; \text{ad } P) &\rightarrow \Omega^0(X; \text{ad } P) \oplus \Omega^1(X; \text{ad } P), \\ D_{A,f}(b, c) &= (d_A^* c, *d_{A,f} c + d_A b) \end{aligned}$$

is a self-adjoint operator called the *perturbed twisted odd signature operator*.

The operator $*Q_{A,f}$ is independent of the choice of Riemannian metric on X since the definition of $\text{Hess}(h_f)_A(b, c)$ does not use the metric, and since

$$\text{Hess}(h_f)_A(b, c) = \langle Q_{A,f}(b), c \rangle = - \int_X \text{tr}(c \wedge *Q_{A,f}(b)).$$

It follows that $d_{A,f}$ is metric independent. In particular, if A is *f-perturbed flat*, then the cohomology of (2.7), and hence the dimension of the kernel of $D_{A,f}$, is independent of the metric on X .

We will give a proof in Lemma 2.3 of Taubes' observation that for any A and f , the operator $D_{A,f}$ is Fredholm and in fact a compact perturbation of D_A . Assuming this, we conclude the following.

Lemma 2.1. *Suppose that f_0 and f_1 are perturbations, A_0 and A_1 are connections so that A_0 is f_0 perturbed flat and A_1 is f_1 perturbed flat. If f_t is a path of perturbations from f_0 to f_1 and A_t is a path of connections from A_0 to A_1 then the spectral flow $\text{SF}(D_{A_t, f_t})$ is independent of the choice of paths A_t and f_t and independent of the choice of Riemannian metric.*

Proof. Since the space of connections and the space of perturbations are contractible and the spectral flow is a homotopy invariant, $\text{SF}(D_{A_t, f_t})$ is independent of the choice of paths A_t from A_0 to A_1 and path f_t from f_0 to f_1 . Since the dimensions of the kernels of D_{A_0, f_0} and D_{A_1, f_1} are independent of the Riemannian metric, it similarly follows that $\text{SF}(D_{A_t, f_t})$ is independent of the choice of Riemannian metric. $\square$

2.5. Locality of $Q_{A,f}$. The operator $D_{A,f}$ is not a differential operator, but it has the property that it differs from a differential operator by an operator which is “localized” in the neighborhood of the solid tori used to construct the perturbations.

Write

$$D_{A,f}(b, c) = D_A(b, c) + (0, -4\pi^2 Q_{A,f}(c))$$

so that $D_{A,f}$ differs from the Dirac operator D_A by $-4\pi^2 Q_{A,f}$.

Proposition 2.2. *Let $S \subset X$ denote the union of the images of the embeddings γ_i , $S = \cup_i \gamma_i(D^2 \times S^1)$.*

- (1) *If $b, b' \in \Omega^1(X; \text{ad } P)$ and $b|_S = b'|_S$, then $Q_{A,f}(b) = Q_{A,f}(b')$.*
- (2) *For any $b \in \Omega^1(X; \text{ad } P)$, $Q_{A,f}(b)$ vanishes outside of S .*
- (3) *If A, A' are connections and $A|_S = A'|_S$, then $Q_{A,f}(b) = Q_{A',f}(b)$ for all $b \in \Omega^1(X; \text{ad } P)$.*

Proof. Suppose that $b \in \Omega^1(X; \text{ad } P)$ vanishes on S . Then for each $c \in \Omega^1(X; \text{ad } P)$,

$$\langle Q_{A,f}(b), c \rangle = \text{Hess}(h_f)_A(b, c) = \frac{\partial^2}{\partial s \partial t} \Big|_{s=t=0} \int_{D^2} f(\text{Hol}(A + sb + tc, x)) \eta(x) d^2x = 0,$$

since the parallel lifts of the loops $\gamma_i(x, u)$, $u \in [0, 2\pi]$ with respect to the connection $A + sb + tc$ are independent of s . Hence $Q_{A,f}(b) = 0$. Part (1) now follows since $Q_{A,f}$ is linear and $b - b'$ vanishes on S .

For part (2), Let $b \in \Omega^1(X; \text{ad } P)$ be arbitrary. Let $U \subset X \setminus S$ and choose $c \in \Omega^1(X; \text{ad } P)$ supported in U . Then

$$\langle Q_{A,f}(b), c \rangle = \langle b, Q_{A,f}(c) \rangle = 0$$

by the same argument given for part (1) since c vanishes on S . Since the L^2 inner product of $Q_{A,f}(b)$ with c vanishes for every c supported on U , $Q_{A,f}(b)$ must vanish on U .

Part (3) is clear. $\square$

Proposition 2.2 implies that the restriction of $D_{A,f}$ to a codimension 0 submanifold whose boundary misses $S = \cup_i \text{Image}(\gamma_i)$ is well-defined. To be precise, suppose that $Y \subset X$ is a compact codimension 0 submanifold. We consider two cases: $S \subset \text{interior}(Y)$ and $S \subset X \setminus Y$.

If $S \subset \text{interior}(Y)$, A is any connection on Y and c is a 1-form on Y with values in $\text{ad } P|_Y$, extend A and c arbitrarily to $\tilde{A}, \tilde{c}$ on X . Then $Q_{\tilde{A},f}(\tilde{c})$ vanishes outside of S and is independent of the choice of extensions. Thus $Q_{A,f}$ is well-defined as an operator on Y . Since D_A is a differential operator it is locally defined and hence has a well defined restriction to Y . It follows that $D_{A,f} = D_A - 4\pi^2 Q_{A,f}$ is well defined on Y .

If $Y \subset X \setminus S$, then a similar argument shows that $D_{A,f}$ restricts to the differential operator D_A on Y .

2.6. An estimate. We next give an argument to show that $D_{A,f}$ differs from D_A by an operator which is bounded on L^2 . This result is essentially contained in Taubes' article [29]. See also [3, Proposition 2.8].

Lemma 2.3. *There exists a constant C so that for all $A \in \mathcal{A}$, $f \in \mathcal{P}$, and $b \in \Omega^1(X; \text{ad } P)$*

$$\|Q_{A,f}(b)\|_{L^2} \leq C\|f\|_{C^2}\|b\|_{L^2}.$$

In particular, $D_{A,f}$ is a compact perturbation of D_A , viewed as operators $L_1^2 \rightarrow L^2$.

Proof. Denote $\text{Hol}(A + sb + tc, x)$ by $H_x(s, t)$, so $H_x : \mathbb{R}^2 \rightarrow SU(n)^N$, and let $f : SU(n)^N \rightarrow \mathbb{R}$ be invariant under the conjugation action of $SU(n)$. Then

$$\text{Hess}(h_f)_A(b, c) = \left. \frac{\partial^2}{\partial s \partial t} \right|_{s=t=0} \int_{D^2} f(H_x(s, t)) \eta(x) d^2x = \int_{D^2} \left. \frac{\partial^2}{\partial s \partial t} \right|_{s=t=0} f(H_x(s, t)) \eta(x) d^2x.$$

We may think of $H_x : \mathbb{R}^2 \rightarrow SU(n)^N \subset \mathbb{C}^{n^2N}$ and extend f to a smooth function $\mathbb{C}^{n^2N} \rightarrow \mathbb{R}$. By the chain and the product rule for vector valued functions we have

$$\begin{aligned} & \left. \frac{\partial^2}{\partial s \partial t} \right|_{s=t=0} f(H_x(s, t)) \\ &= D_{H_x(0,0)} f \circ \left. \frac{\partial^2}{\partial s \partial t} \right|_{s=t=0} H_x(s, t) + \left(\left. \frac{\partial}{\partial s} \right|_{s=0} H_x(s, 0) \right)^T \circ D_{H_x(0,0)}^2 f \circ \left. \frac{\partial}{\partial t} \right|_{t=0} H_x(0, t) \end{aligned}$$

Then

$$\begin{aligned} & \left| \left. \frac{\partial^2}{\partial s \partial t} \right|_{s=t=0} f(H_x(s, t)) \right| \\ & \leq \|D_{H_x(0,0)} f\| \cdot \left\| \left. \frac{\partial^2}{\partial s \partial t} \right|_{s=t=0} H_x(s, t) \right\| + \left\| \left. \frac{\partial}{\partial s} \right|_{s=0} H_x(s, 0) \right\| \cdot \|D_{H_x(0,0)}^2 f\| \cdot \left\| \left. \frac{\partial}{\partial t} \right|_{t=0} H_x(0, t) \right\| \end{aligned}$$

Given $c \in \Omega^1(X; \text{ad } P)$, $i \in \{1, \dots, N\}$, and $x \in D^2$, let $\tilde{c}_{i,x} : [0, 1] \rightarrow su(n)$ be defined as follows. Trivialize $\text{ad } P$ over the disc $D^2 \subset X$, so $\text{ad } P|_{D^2} \cong D^2 \times su(n)$. Parallel translation by A along the path $g_{i,x} : [0, 1] \rightarrow X, t \mapsto \gamma_i(x, e^{2\pi i t})$ trivializes $g_{i,x}^*(\text{ad } P)$ over $[0, 1]$, and this way the pullback $g_{i,x}^*(c)$ is viewed as a 1-form with $su(n)$ coefficients. Contracting with the unit speed vector field $\frac{\partial}{\partial t}$ yields the function $\tilde{c}_{i,x} : [0, 1] \rightarrow su(n)$, i.e. $\tilde{c}_{i,x} = g_{i,x}^*(c)(\frac{\partial}{\partial t})$.

We have the estimate

$$\int_{D^2} \int_0^1 |\tilde{c}_{i,x}(\nu)| \eta(x) d\nu d^2x \leq \int_{D^2} \int_0^1 |c(\nu)| \eta(x) d\nu d^2x \leq C_1 \|c\|_{L^1(S^1 \times D^2)} \leq C \|c\|_{L^1(X)}$$

for some constants C_1, C independent of c .

Given a connection B let $h_{i,x}(B)$ denote the holonomy of B around the loop $\gamma_i(x, -)$. Thus

$$H_x(s, t) = (h_{1,x}(A + sb + tc), \dots, h_{1,x}(A + sb + tc)).$$

Lemma 2.6 of [3] establishes the formulas:

$$\begin{aligned} \left. \frac{d}{dt} \right|_{t=0} h_{i,x}(A + tb) &= h_{i,x}(A) \int_0^1 \tilde{b}_{i,x}(\nu) d\nu \\ \left. \frac{\partial^2}{\partial s \partial t} \right|_{s=t=0} h_{i,x}(A + sb + tc) &= h_{i,x}(A) \int_0^1 \int_0^\nu (\tilde{b}_{i,x}(\nu) \tilde{c}_{i,x}(\mu) + \tilde{c}_{i,x}(\nu) \tilde{b}_{i,x}(\mu)) d\mu d\nu. \end{aligned}$$

Notice that $h_{i,x}(A) \in SU(n)$, and so is bounded independently of A . Thus (letting C denote possibly different constants)

$$\int_{D^2} \left| \left. \frac{d}{dt} \right|_{t=0} H_x(s, 0) \right| \eta(x) d^2x \leq \sum_{i=1}^N C \|b\|_{L^1(X)} = NC \|b\|_{L^1(X)} \leq C \|b\|_{L^2(X)}.$$

The last inequality follows from Cauchy-Schwarz.

Similarly

$$\begin{aligned}
& \int_{D^2} \left| \frac{\partial^2}{\partial s \partial t} \right|_{s=t=0} H_x(s, t) | \eta(x) | d^2 x \\
& \leq C \int_{D^2} \int_0^1 |b(\nu)| d\nu \int_0^1 |c(\mu)| d\mu d^2 x \leq C \left\| \int_0^1 |b(\nu)| d\nu \right\|_{L^2(D^2)} \cdot \left\| \int_0^1 |c(\mu)| d\mu \right\|_{L^2(D^2)} \\
& \leq C \|b\|_{L^2(S^1 \times D^2)} \cdot \|c\|_{L^2(S^1 \times D^2)} \leq C \|b\|_{L^2(X)} \cdot \|c\|_{L^2(X)}.
\end{aligned}$$

Also

$$\|D_{H_x(0,0)} f\| + \|D_{H_x(0,0)}^2 f\| \leq C \|f\|_{C^2}.$$

Thus for any $b, c \in \Omega^1(X; \text{ad } P)$,

$$|\text{Hess}(h_f)_A(b, c)| \leq C \|f\|_{C^2} \|b\|_{L^2} \|c\|_{L^2},$$

and hence $\|Q_{A,f}\| \leq C \|f\|_{C^2}$.

For the last statement of Lemma 2.3, the difference $D_{A,f} - D_A : L_1^2 \rightarrow L^2$ equals the restriction of $-4\pi^2 Q_{A,f}$ to L_1^2 . This restriction can be expressed as the composite of the compact inclusion $L_1^2 \subset L^2$ and the bounded operator $-4\pi^2 Q_{A,f}$, and hence is compact since the compact operators form a 2-sided ideal. $\square$

3. THE POISSON OPERATOR AND THE CALDERÓN PROJECTOR FOR PERTURBED DIRAC OPERATORS ON MANIFOLDS WITH BOUNDARY

A compact manifold with non-empty boundary always embeds as a codimension 0 submanifold of a closed manifold (e.g. its double) in such a way that a given principal bundle and $SU(n)$ connection extend. It follows from Proposition 2.2 that given:

- (1) a 3-manifold with boundary X ,
- (2) a principal bundle $P \rightarrow X$,
- (3) a connection A on P ,
- (4) a collection of embeddings $\gamma_i : D^2 \times S^1 \rightarrow \text{interior}(X)$, and
- (5) a function $f \in \mathcal{P}$

one can unambiguously define operators $Q_{A,f}$ and $D_{A,f}$ on X . In this context we will assume that the connection A is in cylindrical form near the boundary. This means that in a collar neighborhood $[0, \epsilon) \times \partial X$ of ∂X , A is the pullback of a connection a on ∂X with respect to the projection $[0, \epsilon) \times \partial X \rightarrow \partial X$ to the second factor. Then the operator $D_{A,f}$ equals D_A on the collar, and this implies that $D_{A,f}$ takes the Atiyah–Patodi–Singer form (for details see [5, Definitions 2.2 and 2.3])

$$(3.1) \quad D_{A,f}|_{[0,\epsilon) \times \partial X} = D_A|_{[0,\epsilon) \times \partial X} = \gamma\left(\frac{\partial}{\partial u} + S_a\right).$$

Here $S_a : \Omega^*(\partial X; \text{ad } P) \rightarrow \Omega^*(\partial X; \text{ad } P)$ is the tangential operator, given by the formula

$$(3.2) \quad S_a(\alpha_0, \alpha_1, \alpha_2) = (*d_a \alpha_1, -*d_a \alpha_0 - d_a * \alpha_2, d_a * \alpha_1)$$

and γ is the bundle isomorphism given by the formula

$$(3.3) \quad \gamma(\alpha_0, \alpha_1, \alpha_2) = (- * \alpha_2, * \alpha_1, * \alpha_0)$$

(in the formulas (3.3) and (3.2), $*$ refers to the Hodge $*$ operator for the metric on ∂X). The bundle isomorphism γ satisfies $\gamma^2 = -\text{Id}$.

3.1. The invertible double. So far we have shown that $D_{A,f}$ is the sum of a Dirac operator D_A and an L^2 -bounded operator $-4\pi^2 Q_{A,f}$. The operator $Q_{A,f}$ is not a differential operator but it has certain locality properties as explained in Section 2.5. On a manifold with boundary we arranged so that $D_{A,f}$ has a product form in a collar of the boundary.

In the study of Dirac operators on manifolds with boundary, their Calderón projector and their Poisson operator play a central role (cf. [10]). However, since $Q_{A,f}$ is far from being a differential operator, existence of the Calderón projector and of the Poisson operator is a priori not clear. We

therefore treat the existence question and related issues. For the convenience of the reader our presentation will be more or less self-contained.

Let X be a compact connected Riemannian manifold with boundary $\partial X = \Sigma$ and D a self-adjoint Dirac type operator on the Hermitian vector bundle $E \rightarrow X$. We assume that we are in a product situation, i.e. that there is a collar $U = [0, \epsilon) \times \Sigma$ of Σ in which D takes the form

$$(3.4) \quad D = \gamma \left(\frac{d}{dx} + B \right),$$

where γ is a bundle endomorphism and B is a first order self-adjoint elliptic differential operator on $E|_\Sigma$. One has the fundamental relations

$$(3.5) \quad \gamma^2 = -\text{Id}, \quad \gamma^* = -\gamma, \quad \text{and} \quad B\gamma = -\gamma B.$$

Note that D_A is of this form (3.1).

To model $Q_{A,f}$ we assume abstractly that we are given an auxiliary self-adjoint bounded linear operator V in $L^2(X, E)$. In most cases V will be a pseudo-differential operator of order 0, but this is not assumed here. More importantly, we assume the following locality property (cf. Sec. 2.5).

$$(3.6) \quad \text{There is a compact domain } S \subset X \setminus [0, \epsilon) \times \Sigma \text{ such that } M_\varphi V = 0 \text{ for all } \varphi \in C_0^\infty(X \setminus S).$$

Here M_φ denotes the operator of multiplication by φ . Note that by self-adjointness we automatically also have

$$(3.7) \quad VM_\varphi = 0 \quad \text{for} \quad \varphi \in C_0^\infty(X \setminus S).$$

Moreover, if $f, g \in L^2(E)$ such that $f|_W = g|_W$ in a neighborhood W of S then choosing $\psi \in C_0^\infty(W)$ with $\psi \equiv 1$ in a neighborhood of S we find

$$(3.8) \quad V(f - g) = V((1 - \psi)(f - g)) = (VM_{1-\psi})(f - g) = 0.$$

The property (3.6) implies that the restriction of $D + V$ to the collar U equals D and hence takes the product form (3.4). This and self-adjointness is all that is required to prove Green's formula (see [10])

$$(3.9) \quad \langle (D + V)f, g \rangle_{L^2(X, E)} - \langle f, (D + V)g \rangle_{L^2(X, E)} = \langle f|_\Sigma, \gamma(g|_\Sigma) \rangle_{L^2(\Sigma, E)}.$$

An important tool for studying elliptic boundary problems is the *invertible double construction* and the associated Poisson operator. Since it will be important in the sequel we recall the main facts (cf. [10]).

Let $\tilde{X} := X_+ \cup_\Sigma X_-$ where $X_\pm = X$. Let $\tilde{E} \rightarrow \tilde{X}$ be the vector bundle obtained by gluing E with gluing function γ over Σ . More precisely, consider the bundle $E \times \{\pm\} \rightarrow X \times \{\pm\}$. Then

$$\tilde{X} = X \times \{\pm\} / \sim, \quad \text{where } (x, +) \sim (y, -) \text{ if and only if } x = y \text{ and } x \in \Sigma.$$

We write $X_\pm := X \times \{\pm\}$. Also,

$$\tilde{E} = E \times \{\pm\} / \sim \quad \text{where } (x, v, +) \sim (y, w, -) \text{ if and only if } x = y \in \Sigma \text{ and } \gamma(v) = w.$$

Denote by $m : \tilde{X} \rightarrow \tilde{X}$ the reflection interchanging X_+ and X_- , i.e. $m(x, \pm) = (x, \mp)$. Then m is covered by a bundle map $\tilde{m} : \tilde{E} \rightarrow \tilde{E}$ defined by $\tilde{m} : (x, v, \pm) = (x, \mp v, \mp)$. Notice that the restriction of $\tilde{m}$ to $\tilde{E}|_\Sigma$ equals γ since if $x \in \Sigma$, $\tilde{m} : (x, v, \pm) = (x, \mp v, \mp) \sim (x, \gamma(v), \pm)$. The map $\tilde{m}$ preserves the Hermitian metric on E and $\tilde{m}^2 = -\text{Id}$.

The maps m and $\tilde{m}$ induce a unitary map

$$(3.10) \quad \Phi : L^2(\tilde{X}, \tilde{E}) \rightarrow L^2(\tilde{X}, \tilde{E}), \quad \Phi f(x) := \tilde{m}(f(m(x)))$$

satisfying $\Phi^2 = -\text{Id}$. Thus Φ is defined by the diagram

$$\begin{array}{ccc} \tilde{E} & \xleftarrow{\tilde{m}} & \tilde{E} \\ \uparrow \Phi f & & \uparrow f \\ \tilde{X} & \xrightarrow{m} & \tilde{X} \end{array}$$

Then D extends naturally to a Dirac operator $\tilde{D}$ on $\tilde{E}$ satisfying

$$(3.11) \quad \Phi^* \tilde{D} \Phi = -\tilde{D}.$$

Similarly, in view of (3.6), V extends to a bounded operator $\tilde{V}$ satisfying $\Phi^* \tilde{V} \Phi = -\tilde{V}$ which has the same locality properties as V with $\tilde{S} = S \cup m(S)$ instead of S .

Alternatively, we may view $\tilde{D}$ as a self-adjoint realization of a well-posed boundary value problem of $D \oplus (-D)$ acting on the bundle $E \oplus E$ over X . Namely, putting

$$(3.12) \quad \begin{aligned} \Psi : L^2(\tilde{X}, \tilde{E}) &\longrightarrow L^2(X, E \oplus E) \\ \Psi f &:= f|_{X_+} \oplus (\Phi f)|_{X_+} \end{aligned}$$

we find

$$(3.13) \quad \Psi \tilde{D} \Psi^{-1} = D \oplus (-D)$$

and

$$(3.14) \quad \Psi(\text{dom } \tilde{D}) = \Psi(L_1^2(\tilde{E})) = \{f \in L_1^2(X, E \oplus E) \mid (f|_\Sigma)_2 = \gamma(f|_\Sigma)_1\}.$$

It is indeed not difficult to show that $(f|_\Sigma)_2 = \gamma(f|_\Sigma)_1$ is a well-posed boundary condition for $D \oplus (-D)$. The boundary operator of this boundary condition, an orthogonal projection in the pseudodifferential Grassmannian (see [10]), is given by

$$(3.15) \quad \frac{1}{2} \begin{pmatrix} \text{Id} & \gamma \\ -\gamma & \text{Id} \end{pmatrix}.$$

We introduce the usual notations for restriction, extension, and trace operators:

Definition 3.1. Let $r^\pm$ be restriction of sections on $\tilde{X}$ to $X_\pm$ and $e^\pm$ extension by 0 from $X_\pm$. $\varrho^\pm, \varrho$ denote the trace maps

$$(3.16) \quad \begin{aligned} L_s^2(X_\pm, E) &\longrightarrow L_{s-1/2}^2(\Sigma, E), \\ L_s^2(\tilde{X}, \tilde{E}) &\longrightarrow L_{s-1/2}^2(\Sigma, E), \end{aligned} \quad s > 1/2.$$

Similarly, for $0 < |t| < \epsilon$ we denote by ϱ_t the trace map which restricts sections to $\{t\} \times \Sigma$. For ϱ_t (3.16) holds accordingly.

Note that, by the locality property (3.6) of V , we have $r^+ \tilde{V} = V r^+$.

For convenience we abbreviate $T := D + V$ resp. $\tilde{T} := \tilde{D} + \tilde{V}$. The pseudolocality property of $\tilde{V}$ implies that $\tilde{T}$ has well-defined restrictions to $X_\pm$. The following fact is well-known:

Lemma 3.2. *For arbitrary $s \in [0, \infty)$ the trace map ϱ^+ extends to a bounded linear map*

$$\ker T \cap L_s^2(X, E) \rightarrow L_{s-1/2}^2(\Sigma, E).$$

More precisely, if $f \in \ker T \cap L_s^2(X, E)$ then $\varrho^+(f) = \lim_{t \rightarrow 0^+} \varrho_t(f)$ exists in $L_{s-1/2}^2(\Sigma, E)$ and the so defined ϱ^+ is bounded.

With some care the lemma could even be stated for all $s \in \mathbb{R}$. However, Sobolev spaces of negative order on manifolds with boundary are a nuisance. Since we will not need them we content ourselves to the case $s \geq 0$.

Proof. For $V = 0$ the Lemma is well-known ([10, Thm. 13.1], cf. also [23, Chap. I]). If $Tf = 0$ then $Df = 0$ in a collar of Σ . Since the result is local in a collar of Σ , we reach the conclusion. $\square$

In light of Lemma 3.2 we can safely introduce the following notation.

Definition 3.3. Set

$$\begin{aligned} Z_{\pm}^s &:= \{f \in L_s^2(X_{\pm}, E) \mid \tilde{T}|_{X_{\pm}} f = 0\}, \quad s \geq 0, \\ \Lambda_{\pm}^s &:= \varrho^{\pm} Z_{\pm}^{s+1/2}, \quad s \geq -1/2, \\ Z_0 &:= \{f \in L_1^2(\tilde{X}, \tilde{E}) \mid \tilde{T}f = 0\}. \end{aligned}$$

Note that from elliptic regularity we immediately conclude that if $f \in L^2(\tilde{X}, \tilde{E})$ and $\tilde{T}f = 0$ then $f \in L_1^2(\tilde{X}, \tilde{E})$. Since V is only assumed to be L^2 -bounded, this cannot be improved. Also by elliptic theory, $\tilde{T}$ is a Fredholm operator (it is a relatively compact perturbation of $\tilde{D}$) and hence $\dim Z_0 < \infty$.

Proposition 3.4. *If $f \in Z_0$ then f vanishes in a collar of Σ and hence $Z_0 = r^+ Z_0 \oplus r^- Z_0$.*

In particular, $\tilde{D}$ is an invertible operator, called the invertible double of D .

Proof. We apply Green's formula and find

$$\begin{aligned} (3.17) \quad \|\varrho f\|^2 &= -\langle \varrho^+ r^+ f, \gamma \gamma \varrho^+ r^+ f \rangle = -\langle \varrho^+ r^+ f, \gamma \varrho^+ r^+ (\Phi f) \rangle \\ &= -\langle Tf, \Phi f \rangle_{L^2(X_+, E)} + \langle f, T\Phi f \rangle_{L^2(X_+, E)} = 0 \end{aligned}$$

since $f \in Z_0$.

Thus we have $\varrho f = 0$. Since the Dirac operator has the weak unique continuation property (cf. [7]) this implies that f vanishes in a collar neighborhood of Σ . If $V = 0$ then the weak unique continuation property and the connectedness of X imply together with the proven vanishing statement for $f \in Z_0$ that $\tilde{D}$ is invertible. $\square$

Definition 3.5. We say that $D + V$ has the *weak unique continuation property with respect to ∂X* if $f = 0$ is the only element of $Z_+^{1/2}$ with $\varrho^+ f = 0$.

In general we cannot expect $D + V$ to have the weak unique continuation property. However, for small V it is true:

Proposition 3.6. *The operator $D + V$ has the weak unique continuation property with respect to ∂X if and only if $\tilde{D} + \tilde{V}$ is invertible. This is in particular the case if $\|\tilde{D}^{-1}\tilde{V}\| < 1$.*

Remark 3.7. Booss-Bavnbek, Marcolli, and Wang [9] prove the weak unique continuation property for a class of perturbed Dirac operators (perturbations may be nonlinear) arising in Seiberg–Witten theory. However, our perturbations are in general not admissible in their sense (cf. [9, Def. 2.5]). An operator $\mathfrak{P}$ is admissible in the sense of [9] if there is a locally bounded function P such that $|\mathfrak{P}u(x)| \leq P(u, x)|u(x)|$. This implies in particular that $\mathfrak{P}u(x) = 0$ if $u(x) = 0$, i.e. $\mathfrak{P}$ is a local operator. If $\mathfrak{P}$ is a linear operator on smooth sections then this locality automatically implies that $\mathfrak{P}$ is a differential operator of order 0 (i.e. a bundle endomorphism). So the point in [9] is that $\mathfrak{P}$ is allowed to be nonlinear.

Proof of Proposition 3.6. In view of (3.11) and Prop. 3.4 the weak unique continuation property with respect to ∂X implies $Z_0 = 0$ and hence the invertibility of $\tilde{D} + \tilde{V}$.

Conversely, assume that $\tilde{D} + \tilde{V}$ is invertible and consider $f \in Z_+^{1/2}$ with $\varrho^+ f = 0$. Then $e^+ f \in Z_0 = \{0\}$ and hence $f = 0$.

$\|\tilde{D}^{-1}\tilde{V}\| < 1$ implies the invertibility of $\tilde{D} + \tilde{V}$ by means of the Neumann series. $\square$

3.2. Existence of Poisson operator and Calderón projector. Let P_{Z_0} be the orthogonal projection onto Z_0 and denote by ϱ^* the L^2 -dual of ϱ . In other words, for a distributional section f of $\tilde{E}|\Sigma$ and a test function $\varphi \in C^\infty(\tilde{X}, \tilde{E})$ one has $(\varrho^* f, \varphi) := (f, \varrho \varphi)$. In view of Proposition 3.4 one infers from this formula immediately that $P_{Z_0} \varrho^* = 0$.

Furthermore, denote by $\tilde{G}$ be the pseudoinverse of $\tilde{T}$, i.e.

$$(3.18) \quad \tilde{G}f := (\text{Id} - P_{Z_0})(\tilde{T} + P_{Z_0})^{-1}f = \begin{cases} \tilde{T}^{-1}f, & f \in Z_0^\perp, \\ 0, & f \in Z_0. \end{cases}$$

Note that since $\tilde{T}$ maps $L_s^2 \rightarrow L_{s-1}^2$, $0 \leq s \leq 1$ the pseudoinverse maps $L_s^2 \rightarrow L_{s+1}^2$, $-1 \leq s \leq 0$. Together with the mapping properties of the trace map we infer that $\tilde{G}\varrho^*$ maps $L_s^2(\Sigma, E)$ continuously into $L_{s+1/2}^2(\tilde{X}, \tilde{E})$ for each $-1/2 \leq s < 0$. It is a fundamental fact that for $r^\pm \tilde{D}^{-1} \varrho^*$ this can be improved. Namely, one has:

Theorem 3.8. $r^\pm \tilde{D}^{-1} \varrho^*$ maps $L_s^2(\Sigma, E)$ continuously into $L_{s+1/2}^2(X_\pm, E)$ for all $s \geq -1/2$.

For a proof see [10]. It will also follow from our discussion of the continuous dependence of the Calderón projector below. First we want to show that the previous theorem easily extends to hold for $\tilde{T}$ and hence a Calderón projector can be constructed for $\tilde{T}$.

Theorem 3.9. $r^\pm \tilde{G}\varrho^*$ maps $L_s^2(\Sigma, E)$ continuously into $Z_\pm^{s+1/2}$ for $-1/2 \leq s \leq 1/2$.

Furthermore, we have the following resolvent identity relating $\tilde{G}$ and $\tilde{D}^{-1}$:

$$\tilde{G} = \tilde{D}^{-1} - \tilde{G}\tilde{V}\tilde{D}^{-1} - P_{Z_0}\tilde{D}^{-1}.$$

Proof. The resolvent identity follows from

$$(3.19) \quad \begin{aligned} \tilde{G}\tilde{V}\tilde{D}^{-1} &= \tilde{G}(\tilde{T} - \tilde{D})\tilde{D}^{-1} \\ &= (\text{Id} - P_{Z_0})\tilde{D}^{-1} - \tilde{G}. \end{aligned}$$

The operator $P_{Z_0}\tilde{D}^{-1}$ is bounded from $L_s^2(\tilde{X}, \tilde{E})$ to $L_1^2(\tilde{X}, \tilde{E})$ for $s \geq -1$ (see Prop. 3.4 and the comment preceding it). Moreover, $\tilde{G}\tilde{V}\tilde{D}^{-1}$ maps $L_{-1}^2(\tilde{X}, \tilde{E})$ continuously into $L_1^2(\tilde{X}, \tilde{E})$. From Theorem 3.8 and the resolvent identity we thus conclude that $r^\pm \tilde{G}\varrho^*$ maps $L_s^2(\Sigma, E)$ continuously into $L_{s+1/2}^2(X_\pm, E)$.

Moreover, since $\tilde{T}\tilde{G}\varrho^* = (\text{Id} - P_{Z_0})\varrho^* = \varrho^*$ and ϱ^* is supported on Σ , we find that the operator actually maps into $Z_\pm^{s+1/2}$. $\square$

The restriction $-1/2 \leq s \leq 1/2$ in Theorem 3.9 is due to the low regularity assumptions on V . If V is a pseudodifferential operator, then Theorem 3.8 holds verbatim for $\tilde{T}$ instead of $\tilde{D}$. The previous proof can easily be modified to this case.

With Theorem 3.9 and Lemma 3.2 in place we can make the following definitions.

Definition 3.10. (1) Define

$$K_\pm := K(T)_\pm := \pm r^\pm \tilde{G}\varrho^* \gamma : L_s^2(\Sigma, E) \rightarrow Z_\pm^{s+1/2}, \quad -1/2 \leq s \leq 1/2.$$

Then $K := K_+$ is called the *Poisson operator* for $T = D + V$.

(2) Define the projection

$$C_\pm := C_\pm(T) := \varrho^\pm K(T)_\pm : L_s^2(\Sigma, E) \rightarrow L_s^2(\Sigma, E), \quad -1/2 \leq s \leq 1/2.$$

Then $C := C_+$ is called the *Calderón projector* of T .

(3) Taking $s = 0$, the subspace

$$\Lambda := \Lambda(T) := \Lambda_+^0(T) = \text{range } C_+(T) \subset L^2(\Sigma, E)$$

is called the *Cauchy data space* of T .

Proposition 3.11. $C_{\pm}(D+V)$ are complementary orthogonal projections in $L^2(\Sigma, E)$ with $\text{range } C_{\pm} = \Lambda_{\pm}^0$.

Furthermore, $\gamma C_{\pm} \gamma^* = C_{\mp}$. In other words the subspaces $\Lambda_{\pm}^0$ are Lagrangian subspaces with respect to the symplectic form $\omega(x, y) = \langle x, \gamma y \rangle_{L^2}$.

Recall that a Lagrangian subspace is a subspace $V \subset L^2(\Sigma, E)$ satisfying $\gamma V = V^{\perp}$.

Proof. $K_{\pm}$ maps $L^2(\Sigma, E)$ into $Z_{\pm}^{1/2}$ by Theorem 3.9. Hence in view of Lemma 3.2 we find that $C_{\pm}$ is L^2 -bounded with $\text{range } C_{\pm} \subset \Lambda_{\pm}^0$.

We show $\Lambda_+^0 \perp \Lambda_-^0$: pick $\xi \in \Lambda_+^0, \eta \in \Lambda_-^0$ and choose $f \in Z_+^{1/2}, g \in Z_-^{1/2}$ with $\xi = \varrho^+ f, \eta = \varrho^- g$. Then Green's formula yields

$$\begin{aligned} (3.20) \quad (\xi, \eta) &= (\varrho^+ f, \varrho^- g) = -(\varrho^+ f, \gamma \varrho^+(\Phi g)) \\ &= -(Tf, \Phi g)_{L^2(X_+, E)} + (f, T\Phi g)_{L^2(X_+, E)} = 0. \end{aligned}$$

Hence $\Lambda_+^0 \perp \Lambda_-^0$.

Next we show that $C_+ + C_- = \text{Id}$: pick $\xi \in L^2(\Sigma, E)$ and $f \in C^\infty(\tilde{X}, \tilde{E})$. Green's formula gives

$$\begin{aligned} (3.21) \quad ((C_+ + C_-)\xi, \gamma \varrho f)_{\Sigma} &= (\varrho^+ r^+ \tilde{G} \varrho^* \gamma \xi, \gamma \varrho f)_{\Sigma} - (\varrho^- r^- \tilde{G} \varrho^* \gamma \xi, \gamma \varrho f)_{\Sigma} \\ &= (TK_+ \xi, f)_{X_+} - (\tilde{G} \varrho^* \gamma \xi, Tf)_{X_+} + (TK_- \xi, f)_{X_-} - (\tilde{G} \varrho^* \gamma \xi, \tilde{T}f)_{X_-} \\ &= -(\tilde{G} \varrho^* \gamma \xi, \tilde{T}f)_X = -((\text{Id} - P_{Z_0}) \varrho^* \gamma \xi, f)_X \\ &= (-\gamma \xi, \varrho f)_{\Sigma} = (\xi, \gamma \varrho f)_{\Sigma}. \end{aligned}$$

Since $\gamma \varrho : C^\infty(\tilde{X}, \tilde{E}) \rightarrow C^\infty(\Sigma; E)$ is surjective, $(C_+ + C_-)\xi = \xi$.

In sum, $C_{\pm}$ are L^2 -bounded with orthogonal range $\subset \Lambda_{\pm}^0$ and $C_+ + C_- = \text{Id}$. Hence $C_{\pm}$ must be the orthogonal projections onto $\Lambda_{\pm}^0$.

Finally, consider $\xi = \varrho^+ f \in \Lambda_+^0, f \in Z_+^{1/2}$. Then, by (3.11) we have $\Phi f \in Z_-^{1/2}$ and thus $\gamma \xi = \varrho^-(\Phi f) \in \Lambda_-^0$. Hence γ interchanges Λ_+^0 and Λ_-^0 . Since $\Lambda_+^0 \perp \Lambda_-^0$ the spaces $\Lambda_{\pm}^0$ must be Lagrangian and we are done. $\square$

3.3. Continuous dependence of the Calderón projector. Let us first describe the set-up in which we want to study the dependence of the Calderón projector on its input data: We consider the class $\mathcal{D}$ of Dirac operators on E which are in cylindrical form near Σ and the class $\mathcal{V}$ of auxiliary operators satisfying (3.6). We say that a family $D_t \in \mathcal{D}$ varies continuously if all coefficients of D_t (in any local chart) vary continuously. Moreover, we equip $\mathcal{V}$ with the usual norm topology of $\mathcal{B}(L^2(X, E))$. Finally, we put $(\mathcal{D} \times \mathcal{V})(n) := \{(D, V) \in \mathcal{D} \times \mathcal{V} \mid \dim Z_0(D + V) = n\}$.

We start with some elementary statements on the continuous dependence of P_{Z_0} and $\tilde{G}$:

Proposition 3.12. *The maps*

$$\begin{aligned} (3.22) \quad P_{Z_0}, \tilde{G} : (\mathcal{D} \times \mathcal{V})(n) &\rightarrow \mathcal{B}(L^2(\tilde{X}, \tilde{E}), L_1^2(\tilde{X}, \tilde{E})), \\ P_{Z_0}(D, V) &:= \text{Proj}(\ker(\tilde{D} + \tilde{V})), \\ \tilde{G}(D, V) &:= (\text{Id} - P_{Z_0}(D, V))(\tilde{D} + \tilde{V} + P_{Z_0}(D, V))^{-1}, \end{aligned}$$

are continuous.

Note that this statement is slightly stronger than the continuity of $P_{Z_0}, \tilde{G}$ as maps into $\mathcal{B}(L^2(\tilde{X}, \tilde{E}))$.

Proof. Operators in $\mathcal{D}$ are in cylindrical form near Σ , hence $\mathcal{D} \times \mathcal{V} \rightarrow \mathcal{B}(L_1^2(\tilde{X}, \tilde{E}), L^2(\tilde{X}, \tilde{E})), (D, V) \mapsto \tilde{D} + \tilde{V}$ is continuous. Thus also $\mathbb{C} \times \mathcal{D} \times \mathcal{V} \rightarrow \mathcal{B}(L_1^2(\tilde{X}, \tilde{E}), L^2(\tilde{X}, \tilde{E})), (\lambda, D, V) \mapsto \tilde{D} + \tilde{V} - \lambda$ is continuous. For a fixed pair $(D_0, V_0) \in (\mathcal{D} \times \mathcal{V})$ there is an $\varepsilon > 0$ such that 0 is the only spectral

point of $\tilde{D}_0 + \tilde{V}_0$ in the closed disc $\{z \in \mathbb{C} \mid |z| \leq \varepsilon\}$. By continuity this remains true for (D, V) in a sufficiently small neighborhood of (D_0, V_0) in $(\mathcal{D} \times \mathcal{V})(n)$. For these (D, V) we have

$$(3.23) \quad P_{Z_0}(D, V) = \frac{1}{2\pi i} \oint_{|z|=\varepsilon} (\lambda - \tilde{D} - \tilde{V})^{-1} d\lambda.$$

Since the inversion is continuous this proves the continuity statement on P_{Z_0} .

The continuity statement on $\tilde{G}(D, V)$ now follows immediately from the defining formula (3.22). $\square$

The resolvent identity in Theorem 3.9 shows that for $(D, V) \in (\mathcal{D} \times \mathcal{V})(n)$ the Poisson operator and hence $C_+(D+V)$ depend continuously on V : namely $K_+(D+V) = K_+(D) - r^+(\tilde{G}(D, V)\tilde{V}\tilde{D}^{-1} + P_{Z_0}(D, V)\tilde{D}^{-1})\varrho^*\gamma$ and by the previous Proposition

$$(D, V) \mapsto r^+(\tilde{G}(D, V)\tilde{V}\tilde{D}^{-1} + P_{Z_0}(D, V)\tilde{D}^{-1})\varrho^*\gamma$$

is continuous $(\mathcal{D} \times \mathcal{V})(n) \rightarrow \mathcal{B}(L_{-1/2}^2(\Sigma, E), L_1^2(X_+, E))$. However, the dependence on D itself is more subtle. In order to explore this we have to look a bit closer at the construction of the Calderón projector.

Fix a $c \in \mathbb{R}$ with $c \notin \text{spec } B$. We abbreviate the spectral projections $P_{>c} := 1_{(c, \infty)}(B)$, $P_{\leq c} := 1_{(-\infty, c]}(B)$ etc. Then we set, for $\xi \in L_s^2(\Sigma, E)$,

$$(3.24) \quad Q(B)\xi(x) := e^{-xB}P_{\geq c}\xi, \quad x \geq 0.$$

We first need to study the mapping properties of $Q(B)$ with respect to Sobolev norms. The results we are going to present are fairly standard applications of the spectral theorem. However, to be as self-contained as possible we present the details.

Let $\epsilon > 0$, $\epsilon < 1$ and fix a smooth cut-off function $\varphi : \mathbb{R} \rightarrow [0, 1]$ with $\varphi(x) = 1$ for $|x| \leq \epsilon/2$ and $\varphi(x) = 0$ for $|x| \geq \epsilon$. Let $\psi : \mathbb{R} \rightarrow \mathbb{R}$ be any function in $C_0^\infty((-\epsilon, \epsilon) \setminus \{0\})$, for example $\psi = \varphi'$. In the following proposition we only use the restrictions of φ and ψ to $\{x \geq 0\}$.

Proposition 3.13.

1. For all $s, s' \in \mathbb{R}$ we have $e^+\psi Q(B) \in \mathcal{B}(L_s^2(\Sigma, E), L_{s'}^2(\tilde{X}, \tilde{E}))$.
2. For $-1/2 \leq s \leq 1/2$ the operator $\varphi Q(B)$ maps $L_s^2(\Sigma, E)$ continuously into $L_{s+1/2}^2(X_+, \tilde{E})$.

Proof. First we need to recall the definition of Sobolev spaces on a manifold with boundary. Since B is elliptic the Sobolev norms on $L_s^2(\Sigma, E)$ can be defined using $\text{Id} + |B|$:

$$(3.25) \quad \langle f, g \rangle_{L_s^2} := \langle (\text{Id} + |B|)^s f, (\text{Id} + |B|)^s g \rangle_{L^2}$$

for $f, g \in L_s^2(\Sigma, E)$. Furthermore, for a nonnegative integer $k \in \mathbb{Z}_+$ a Sobolev norm for $L_k^2(\mathbb{R}_+ \times \Sigma, E)$ is given by

$$(3.26) \quad \|f\|_{L_k^2}^2 = \sum_{j=0}^k \int_0^\infty \|\partial_x^j (\text{Id} + |B|)^{k-j} f(x)\|_{L^2(\Sigma, E)}^2 dx.$$

The Sobolev norms on X_+ are obtained by patching together the usual interior Sobolev norms with the norms (3.26).

1. By complex interpolation it suffices to prove the claim for $s' \in \mathbb{Z}_+$. Applying the Leibniz rule to $\partial^j (\text{Id} + |B|)^{k-j} \psi Q(B)$ we see that in view of (3.26) it is in fact sufficient to show that for any $k \in \mathbb{Z}_+$ the operator $(\text{Id} + |B|)^k \psi Q(B)$ is bounded $L_s^2(\Sigma, E) \rightarrow L^2(X_+, E)$.

Furthermore, for $\xi \in L_s^2(\Sigma, E)$ we have $(\text{Id} + |B|)^s \xi \in L^2(\Sigma, E)$ and

$$(\text{Id} + |B|)^k \psi Q(B)\xi = (\text{Id} + |B|)^{k-s} \psi Q(B)((\text{Id} + |B|)^s \xi)$$

so that it suffices to prove the L^2 -boundedness of $(\text{Id} + |B|)^k \psi Q(B)$ for all k .

Next, since $\psi \in C_0^\infty(0, \epsilon)$ there is an $x_0 > 0$ such that $\psi(x) = 0$ for $x \leq x_0$. For $x_0 \leq x \leq \epsilon$ the spectral theorem gives

$$(3.27) \quad \|(\text{Id} + |B|)^k e^{-x_B} P_{\geq c}\|_{L^2(\Sigma, E)} = \sup_{\lambda \geq c} |(1 + \lambda)^k e^{-x\lambda}| \leq C(k, x_0, c, \epsilon)$$

and thus for $\xi \in L^2(\Sigma, E)$

$$(3.28) \quad \begin{aligned} \|\psi(\text{Id} + |B|)^k Q(B)\xi\|_{L^2(X_+, E)}^2 &\leq \|\psi\|_\infty^2 \int_{x_0}^\epsilon \|(\text{Id} + |B|)^k e^{-x_B} P_{\geq c}\xi\|_{L^2(\Sigma, E)}^2 dx \\ &\leq \epsilon \|\psi\|^2 C(k, x_0, c, \epsilon)^2 \|\xi\|^2. \end{aligned}$$

The proof of 1. now follows since ψ has compact support in $(0, \infty)$ and hence in this case extension by 0 is bounded $L_{s'}^2(X_+) \rightarrow L_{s'}^2(\tilde{X})$ for any s' .

2. Let $(E_\lambda)_{\lambda \in \mathbb{R}}$ be the spectral resolution of B . Then we estimate for $\xi \in L^2(\Sigma, E)$

$$(3.29) \quad \begin{aligned} \|\varphi(\text{Id} + |B|)^{1/2} Q(B)\xi\| &\leq \int_0^\infty \|(\text{Id} + |B|)^{1/2} \varphi(x) e^{-x_B} P_{\geq c}(B)\xi\|^2 dx \\ &\leq \int_c^\infty \int_0^\epsilon (1 + |\lambda|) e^{-2x\lambda} dx d\langle E_\lambda \xi, \xi \rangle \\ &\leq C(c, \epsilon) \|\xi\|^2. \end{aligned}$$

For $\xi \in L_{-1/2}^2(\Sigma, E)$ we can now estimate using (3.29)

$$(3.30) \quad \begin{aligned} \|\varphi Q(B)\xi\|_{L^2(X_+, E)} &= \|\varphi(\text{Id} + |B|)^{1/2} Q(B)(\text{Id} + |B|)^{-1/2} \xi\|_{L^2(X_+, E)} \\ &\leq \|\varphi(\text{Id} + |B|)^{1/2} Q(B)\|_{L^2(\Sigma, E) \rightarrow L^2(X_+, E)} \|(\text{Id} + |B|)^{-1/2} \xi\|_{L^2(\Sigma, E)} \end{aligned}$$

proving the continuity of $\varphi Q(B)$ from $L_{-1/2}^2(\Sigma, E)$ to $L^2(X_+, E)$.

Next consider $\xi \in L_{1/2}^2(\Sigma, E)$. Then analogously we find (taking $\psi = \varphi'$ and applying part 1)

$$(3.31) \quad \begin{aligned} \|\partial_x \varphi Q(B)\xi\|_{L^2(X_+, E)} &\leq \|\varphi' Q(B)\xi\|_{L^2(X_+, E)} + \|\varphi B(\text{Id} + |B|)^{-1/2} Q(B)(\text{Id} + |B|)^{1/2} \xi\|_{L^2} \\ &\leq C_1 \|\xi\|_{L^2} + C_2 \|(\text{Id} + |B|)^{1/2} \xi\|_{L^2} \end{aligned}$$

and

$$(3.32) \quad \begin{aligned} \|\varphi(\text{Id} + |B|) Q(B)\xi\|_{L^2(X_+, E)} &= \|\varphi(\text{Id} + |B|)^{1/2} Q(B)(\text{Id} + |B|)^{1/2} \xi\|_{L^2} \\ &\leq C(c, \epsilon) \|(\text{Id} + |B|)^{1/2} \xi\|_{L^2}. \end{aligned}$$

In view of (3.26) this proves the continuity of $\varphi Q(B)$ from $L_{1/2}^2(\Sigma, E)$ to $L_1^2(X_+, E)$.

The complex interpolation method [30, Sec. 4.2] now yields the assertion for all $s \in [-1/2, 1/2]$. $\square$

Proposition 3.14. *Let $\text{Ell}^1(\Sigma, E)$ be the set of first order self-adjoint elliptic differential operators on $E|_\Sigma$. Then for $\psi \in C_0^\infty((-\epsilon, \epsilon) \setminus 0)$ the map*

$$\text{Ell}^1(\Sigma, E)_c := \{B \in \text{Ell}^1(\Sigma, E) \mid c \notin \text{spec } B\} \longrightarrow \mathcal{B}(L^2(\Sigma, E), L^2(X_+, E)), \quad B \mapsto \psi Q(B)$$

is continuous.

Proof. Again let $x_0 > 0$ be such that $\psi(x) = 0$ for $0 \leq x \leq x_0$. By the spectral theorem the map

$$(3.33) \quad [x_0, \epsilon] \times \text{Ell}^1(\Sigma, E)_c \longrightarrow \mathcal{B}(L^2(\Sigma, E)), \quad (x, B) \mapsto e^{-x_B} P_{\geq c}(B)$$

is continuous.

Now fix $B_0 \in \text{Ell}^1(\Sigma, E)_c$ and a $\delta > 0$. By compactness there is a neighborhood W of B_0 such that for $B \in W$ and all $x_0 \leq x \leq \epsilon$

$$(3.34) \quad \|e^{-x_B} P_{\geq c}(B) - e^{-x_{B_0}} P_{\geq c}(B_0)\|_{L^2(\Sigma, E)} \leq \delta / \sqrt{\epsilon}.$$

Thus for $\xi \in L^2(\Sigma, E)$

$$(3.35) \quad \|\psi Q(B)\xi - \psi Q(B_0)\xi\|^2 \leq C^2 \int_{x_0}^{\epsilon} \|e^{-xB} P_{\geq c}(B) - e^{-xB_0} P_{\geq c}(B_0)\|_{L^2(\Sigma, E)}^2 dx \|\xi\|^2 \leq C^2 \delta^2 \|\xi\|^2.$$

Thus $\|\psi Q(B) - \psi Q(B_0)\| \leq C\delta$, completing the proof. $\square$

Let

$$(3.36) \quad R(B)\xi(x) := \begin{cases} -e^{-xB} P_{\geq c}\gamma\xi & \text{for } x \geq 0, \\ e^{-xB} P_{\leq c}\gamma\xi & \text{for } x < 0, \end{cases}$$

and define the *approximate Poisson operator* $K_{\pm}^0 := \pm r^{\pm} \varphi R(B) \gamma$.

Applying Proposition 3.13 and 3.14 (and its obvious counterparts for $x \leq 0$) to $R(B)$ gives:

Proposition 3.15. 1. *If $\psi \in C_0^\infty((-\epsilon, \epsilon) \setminus \{0\})$ then for all $s, s' \in \mathbb{R}$,*

$$\psi R(B) \in \mathcal{B}(L_s^2(\Sigma, E), L_{s'}^2(\tilde{X}, \tilde{E})).$$

Moreover, as an element of $\mathcal{B}(L^2(\Sigma, E), L^2(\tilde{X}, \tilde{E}))$, $\psi R(B)$ depends continuously on the coefficients of B as long as $c \notin \text{spec } B$.

2. *The map $\varphi R(B)$ maps $L^2(\Sigma, E)$ continuously into $L_{1/2}^2(X_- \amalg X_+, \tilde{E})$.*

If $f \in C^\infty(\tilde{X}, \tilde{E})$ then Green's formula gives

$$(3.37) \quad \begin{aligned} (\varphi R\xi, \tilde{T}f)_{\tilde{X}} &= (r^+ \varphi R\xi, r^+ \tilde{T}f)_{X^+} + (r^- \varphi R\xi, r^- \tilde{T}f)_{X^-} \\ &= (-\gamma P_{\geq c}\gamma\xi, \varrho f)_{\Sigma} - (\gamma P_{\leq c}\gamma\xi, \varrho f)_{\Sigma} + (S(B)\xi, f)_{\tilde{X}} \\ &= ((\varrho^* + S(B))\xi, f)_{\tilde{X}} \end{aligned}$$

with $S(B)\xi = \varphi' \gamma R\xi$. In view of Prop. 3.15 we have in particular that $S(B) \in \mathcal{B}(L^2(\Sigma, E), L^2(\tilde{X}, \tilde{E}))$ and that it depends continuously on the coefficients of B . Note furthermore, that $S(B)$ maps in fact into $L_{s, \text{comp}}^2(\tilde{X} \setminus \Sigma, \tilde{E})$ for all $s \geq 0$. Hence we have shown that

$$(3.38) \quad \tilde{T}\varphi R = \varrho^* + S(B).$$

The definition of $K_{\pm}^0$ immediately gives

$$(3.39) \quad \begin{aligned} \varrho^+ K_+^0 \xi &:= \lim_{t \rightarrow 0^+} \varrho_t K_+^0 \xi = P_{\geq c} \xi, \\ \varrho^- K_-^0 \xi &:= \lim_{t \rightarrow 0^-} \varrho_t K_-^0 \xi = P_{\leq c} \xi. \end{aligned}$$

Note that we assumed $c \notin \text{spec } B$.

(3.38) and (3.39) allow to express $K_{\pm}(\tilde{D} + \tilde{V})$ and $C_{\pm}(\tilde{D} + \tilde{V})$ in terms of the operator $R(B)$. Namely, we have in view of (3.38)

$$(3.40) \quad (\text{Id} - P_{Z_0})\varphi R = \tilde{G}\tilde{T}\varphi R = \tilde{G}\varrho^* + \tilde{G}S(B)$$

and hence

$$(3.41) \quad \begin{aligned} K_+ &= r^+ \tilde{G}\varrho^* \gamma \\ &= r^+ (\text{Id} - P_{Z_0})\varphi R \gamma - r^+ \tilde{G}S(B)\gamma \\ &= K_+^0 - r^+ P_{Z_0}\varphi R \gamma - r^+ \tilde{G}S(B)\gamma. \end{aligned}$$

Similarly,

$$(3.42) \quad K_- = K_-^0 + r^- P_{Z_0}\varphi R \gamma + r^- \tilde{G}S(B)\gamma.$$

Finally, this gives the formulas:

$$(3.43) \quad \begin{aligned} C_+ &= P_{\geq c}(B) - \varrho^+ \tilde{G}S(B)\gamma, \\ C_- &= P_{\leq c}(B) + \varrho^- \tilde{G}S(B)\gamma. \end{aligned}$$

In view of Proposition 3.12 and Proposition 3.15 the operator $\varrho^\pm \tilde{G}S(B)\gamma$ maps $L^2(\Sigma, E)$ continuously into $L^2_{1/2}(\Sigma, E)$ and, as long as $\dim Z_0$ is constant, it depends continuously on D and V .

Since $c \notin \text{spec}(B)$ the operators $P_{\geq c}(B), P_{\leq c}(B)$ depend continuously on B . Note that the choice of c is irrelevant. To show the continuity of C_+ at D and V we choose a c not in $\text{spec}(B)$. Then we will have $c \notin \text{spec}(B)$ in a full neighborhood of D and V . Of course, $S(B)$ also depends on c .

Summing up we have proved:

Theorem 3.16. *We consider the class $\mathcal{D}$ of Dirac operators on E which are in cylindrical form near Σ and the class $\mathcal{V}$ of auxiliary operators satisfying (3.6). We say that a family $D_t \in \mathcal{D}$ varies continuously if all coefficients of D_t (in any local chart) vary continuously. Moreover, we equip $\mathcal{V}$ with the usual norm topology of $\mathcal{B}(L^2(X, E))$. Then for each $n \in \mathbb{Z}_+$ the map*

$$\{(D, V) \in \mathcal{D} \times \mathcal{V} \mid \dim Z_0(D + V) = n\} \longrightarrow \mathcal{B}(L^2(\Sigma, E)), \quad (D, V) \mapsto C(D + V)$$

is continuous.

Note also that in view of (3.43) the operator $C - P_{\geq 0}(B)$ maps L^2 continuously into $L^2_{1/2}$, i.e. ‘is an operator of order $-1/2$ ’. If V is a pseudodifferential operator then one easily shows that $\varrho^\pm \tilde{G}S(B)$ is a smoothing operator and we recover, though with a different argument, the well-known fact that $C - P_{\geq 0}(B)$ is smoothing [28, Prop. 2.2], [21, Prop. 4.1].

3.4. Application to the perturbed Dirac operator. We turn to our specific operator $D_{A,f} = D_A - 4\pi^2 Q_{A,f}$. The discussion in Section 2.5 shows that $Q_{A,f}$ satisfies (3.6). Hence the general results of Sections 3.2 and 3.3 apply. In particular there exist the Poisson operator $K_+(D_{A,f})$ and the Calderón projector $P_{A,f} := C_+(D_{A,f})$. Next we show that in a neighborhood of the flat connections and for “small” f the weak unique continuation property with respect to ∂X holds for $D_{A,f}$.

Lemma 3.17. *There is an open neighborhood $U \subset \mathcal{A}$ of the set of flat connections and a constant C (which depends only on X as a Riemannian manifold) so that given any $A \in U$,*

$$\|\tilde{D}_A^{-1}\| \leq C.$$

Proof. Given a flat connection A and a gauge transformation g over X the operators D_A and $D_{g \cdot A}$ are unitarily equivalent (i.e. $D_{g \cdot A}(a) = g D_A g^{-1}(a)$), and this equivalence extends to $\tilde{D}_A$ since $\mathcal{G}$ acts on the Lie algebra coefficients and $\gamma = \pm *$ acts on the differential forms, thus $g_x \gamma_x(v) = \gamma_x(g_x(v))$ for $x \in \Sigma$. Hence the spectra of $\tilde{D}_A$ and $\tilde{D}_{g \cdot A}$ coincide.

The space $\mathcal{M}$ of flat connections modulo gauge transformations on X is compact, being homeomorphic to the compact real algebraic variety $\text{Hom}(\pi_1 X, SU(n))/\text{conjugation}$. Thus there is a lower bound, say k , on the absolute value of the smallest eigenvalue of $\tilde{D}_A$ as A varies in the set of flat connections on X . It follows that the largest eigenvalue of $\tilde{D}_A^{-1}$ is bounded by $1/k$, and hence $\|\tilde{D}_A^{-1}\| \leq 1/k$. Set $C = 2/k$, then by continuity the set $U = \{A \in \mathcal{A} \mid \|\tilde{D}_A^{-1}\| \leq C\}$ is a neighborhood of the set of flat connections. $\square$

Proposition 3.18. *There is an $\epsilon > 0$ so that given any connection A in the neighborhood U of Lemma 3.17, and any $f \in \mathcal{P}$ satisfying $\|f\|_{C^r} < \epsilon$, then $\tilde{D}_A - 4\pi^2 \tilde{Q}_{A,f}$ is invertible and hence has the weak unique continuation property with respect to ∂X .*

Proof. The Neumann series shows that $\tilde{D}_A - 4\pi^2 \tilde{Q}_{A,f}$ is invertible if $\|\tilde{D}_A^{-1} 4\pi^2 \tilde{Q}_{A,f}\| < 1$.

By Lemmas 3.17 and 2.3 there exists a constant C independent of A and f such that

$$\|\tilde{D}_A^{-1} 4\pi^2 \tilde{Q}_{A,f}\| \leq 4\pi^2 \|\tilde{D}_A^{-1}\| \|\tilde{Q}_{A,f}\| \leq C \|f\|_{C^2}.$$

Hence we may choose $\epsilon := \frac{1}{C}$. Proposition 3.6 then shows that $D_{A,f}$ has the weak unique continuation property. $\square$

Theorem 3.19. *The map $U \times B_{\mathcal{P}}(0, \epsilon) \rightarrow \mathcal{B}(L^2(\Sigma, E))$ sending a pair (A, f) to the Calderón projector $C_{A,f}$ of $D_{A,f}$ is continuous, and hence the Cauchy data spaces $\Lambda_{A,f} := \text{range}(C_{A,f})$ vary continuously (in the gap metric on the Grassmannian of Lagrangian subspaces of $L^2(\Sigma, E)$) with respect to A, f .*

Proof. By Prop. 3.18 we have $Z_0(\tilde{D}_A - 4\pi^2 \tilde{Q}_{A,f}) = 0$ for all $(A, f) \in \mathcal{A} \times B_{\mathcal{P}}(0, \epsilon)$. Thus the claim follows from Theorem 3.16. $\square$

4. AN APPLICATION

Theorem 3.19 allows us to use the technology developed in [25] and [12] to study how the spectral flow of a path of operators D_{A_t, f_t} on a closed 3-manifold behaves with respect to a decomposition along a separating surface. In particular, the proof of Nicolaescu's adiabatic limit theorem and Lemma 3.2 of [12], which consider the effect of stretching the collar of X on the Cauchy data spaces, do not depend on the operators being Dirac operators, but only on the fact that they have the Atiyah–Patodi–Singer form on a collar. Thus Theorem 5.1 of [12], and all its consequences (Section 6 of [12]) are true for the operators $D_{A,f}$.

This will be used crucially in forthcoming calculations of the $SU(3)$ Casson invariant for Brieskorn spheres [6]. We finish this article with a useful application of our main result. We outline a different proof of a result of Taubes which is the main ingredient in his proof that his invariant equals Casson's $SU(2)$ invariant for homology 3-spheres, and is the starting point for proofs of various approaches to problems known as the Atiyah–Floer conjecture. It is also one of the simplest non-trivial examples to which this machinery can be applied, since this situation avoids the delicate problem of singularities in the moduli space.

First, we set up some notation. Let X be a homology 3-sphere. Let $P \rightarrow X$ be the trivial principal $SU(2)$ bundle and let $\mathcal{A}(X)$ denote the $(L^2_1 \text{ completion of the})$ space of $SU(2)$ connections on P . Denote by $\mathcal{G}(X)$ the $(L^2_2 \text{ completion of the})$ gauge group $\text{Map}(X, SU(2))$. Then let $\mathcal{A}^*(X)$ be the subspace of $\mathcal{A}(X)$ consisting of irreducible connections, that is, those connections whose stabilizer in $\mathcal{G}(X)$ is equal to the center $\{\pm \text{Id}\} \subset \mathcal{G}(X)$. Thus $\mathcal{A}^*(X)/\mathcal{G}(X)$ is a smooth Hilbert manifold.

Suppose that X is given a Heegard decomposition $X = X_1 \cup_{\Sigma} X_2$ of genus $g > 2$. Choose embeddings γ_i of solid tori so that $S = \cup_i \gamma_i(D^2 \times S^1)$ is contained in the interior of X_2 , and choose $f \in \mathcal{P}$ so that the restriction of $cs + h_f$ to the space $\mathcal{A}^*(X)$ of irreducible connections has only non-degenerate critical points (i.e. the kernel of $D_{A,f}$ is zero at every irreducible perturbed flat connection A). In this context the fact that such f are dense in any neighborhood of $0 \in \mathcal{P}$ is proven in [29] and [17]. We will restrict f further below.

It follows that the critical points of $cs + h_f : \mathcal{A}^*(X)/\mathcal{G}(X) \rightarrow \mathbb{R}/\mathbb{Z}$ are finite and isolated. Let $\mathcal{M}_f^*(X)$ denote the critical set, i.e. the moduli space of irreducible perturbed flat connections on X :

$$\mathcal{M}_f^*(X) = \{A \in \mathcal{A}^*(X) \mid d(cs + h_f)(A) = 0\}/\mathcal{G}(X).$$

To a pair $[A_0], [A_1] \in \mathcal{M}_f^*(X)$, Taubes assigns the sign

$$\text{Sign}_T([A_0], [A_1]) = (-1)^{\text{SF}(D_{A_t, f})}$$

where A_t is any path in $\mathcal{A}^*(X)$ joining lifts of $[A_0]$ and $[A_1]$. In fact an application of the index theorem shows that the mod 8 reduction of $\text{SF}(D_{A_t, f})$ depends only on the gauge equivalence classes $[A_0]$ and $[A_1]$.

Restricting irreducible connections to the pieces of the Heegard decomposition leads to a diagram (with the obvious notation)

$$\begin{array}{ccc}
 & \mathcal{M}^*(X_1) & \\
 \nearrow & & \searrow \\
 \mathcal{M}_f^*(X) & & \mathcal{M}^*(\Sigma) \\
 \searrow & & \nearrow \\
 & \mathcal{M}_f^*(X_2) &
 \end{array}$$

All arrows are embeddings. The space $\mathcal{M}^*(\Sigma)$ is a smooth $6g - 6$ dimensional symplectic manifold. (We will describe the symplectic structure below.) The subspaces $\mathcal{M}^*(X_1)$ and $\mathcal{M}_f^*(X_2)$ are smooth and orientable $3g - 3$ dimensional Lagrangian submanifolds ([29, 18]). They intersect transversely in the compact 0-dimensional manifold $\mathcal{M}_f^*(X)$ for appropriate f , namely those perturbations f so that $cs + h_f$ has only non-degenerate critical points in $\mathcal{A}^*(X)$. Notice that since S lies in the interior of X_2 , perturbed flat connections restrict to flat connections on X_1 and Σ .

We also assume that $\|f\|_{C^2}$ is small enough so the following requirements hold:

- (1) The space $\mathcal{M}_f^*(X_2)$ is diffeomorphic to the flat (i.e. unperturbed) moduli space $\mathcal{M}^*(X_2)$. This is possible since, as f varies, $\mathcal{M}_f^*(X_2)$ varies by a (Legendrian) cobordism ([18]) and so for small enough f the cobordism is a product. (Some care must be taken since $\mathcal{M}_f^*(X_2)$ is non-compact, but the results of [18] can be used to reach this conclusion. Alternatively, Taubes shows that the perturbations can be chosen to fix the reducible stratum.)
- (2) The perturbation should be small enough so that $\mathcal{M}_f^*(X_2)$ lies in the neighborhood U specified by Lemma 3.17. That this is possible follows again from [18].
- (3) The perturbation should be small enough so that the requirement $\|f\| < \epsilon$ of Proposition 3.18 holds on X_2 .

Then Theorem 3.19 implies that the map taking $A \in \mathcal{M}_f^*(X_2)$ to the Cauchy data space $\Lambda_{A,f}$ for $D_{A,f}$ is continuous.

In this language Casson assigns the sign (after orienting the moduli spaces $\mathcal{M}^*(X_1)$, $\mathcal{M}_f^*(X_2)$ and $\mathcal{M}^*(\Sigma)$)

$$\text{Sign}_C([A_0], [A_1]) = \begin{cases} 1 & \text{if } \mathcal{M}^*(X_1) \cdot_{[A_1]} \mathcal{M}_f^*(X_2) = \mathcal{M}^*(X_1) \cdot_{[A_0]} \mathcal{M}_f^*(X_2) \text{ and,} \\ -1 & \text{otherwise,} \end{cases}$$

where $\mathcal{M}^*(X_1) \cdot_{[A_1]} \mathcal{M}_f^*(X_2) \in \{\pm 1\}$ denotes the local transverse intersection number of $\mathcal{M}^*(X_1)$ and $\mathcal{M}_f^*(X_2)$ in $\mathcal{M}^*(\Sigma)$ at $[A_1]$.

Taubes' proof that his invariant equals Casson's invariant (up to an overall sign which depends on a method for fixing particular orientations of all the moduli spaces) reduces to the assertion that $\text{Sign}_T = \text{Sign}_C$. Notice that we are looking at relative signs, so that the validity of $\text{Sign}_T = \text{Sign}_C$ is independent of the choice of orientations of the moduli spaces.

We begin with a lemma whose proof appears to be well known to those who know it well (e.g. [16]), although we were unable to find an explicit statement in the literature. The statement is implicit in [13], but is described in the language of stable holomorphic bundles. We are indebted to G. Daskalopoulos for help with the argument.

Lemma 4.1. *The space $\mathcal{F}^*(\Sigma)$ of irreducible flat $SU(2)$ connections over a closed 2-manifold Σ of genus greater than 2 is simply connected. The moduli space $\mathcal{M}^*(\Sigma)$ of such connections is also simply connected.*

Proof. Before we start the proof, we remark that in the statement of Lemma 4.1, the space $\mathcal{F}^*(\Sigma)$ can mean either the smooth flat irreducible connections, or the L_r^2 completion of this space for any r large enough so that the L_{r+1}^2 completion of the gauge group consists of continuous gauge transformations, e.g. $r > 0$.

We first set up notation. We consider the trivial rank 2 smooth vector bundle $E = \mathbb{C}^2 \times \Sigma \rightarrow \Sigma$ endowed with a Hermitian metric. We fix a holomorphic structure on Σ . The space $\mathcal{A}$ of $SU(2)$ connections on E is identified with the space of holomorphic structures with holomorphically trivial determinant on E via the standard construction (see [1]) since Σ is 1 complex-dimensional and so every structure is integrable. We denote by $\mathcal{A}^s$ the subspace of stable holomorphic structures on E (see [1]). We take the L_r^2 completions of these spaces.

Denote by $\mathcal{F} \subset \mathcal{A}$ the subspace of flat connections, and by $\mathcal{F}^*$ the subspace of irreducible flat connections. We take $\mathcal{G}$ to be the (L_{r+1}^2) completion of the $SU(2)$ gauge group $\text{Map}(\Sigma, SU(2))$.

The slice theorem (see e.g. [15]) shows that the quotient map $\mathcal{A} \rightarrow \mathcal{A}/\mathcal{G}$, when restricted to the irreducible connections $\mathcal{A}^*$ gives a locally trivial map, and hence a fibration $\mathcal{A}^* \rightarrow \mathcal{A}^*/\mathcal{G}$ with fiber $\mathcal{G}/\pm \text{Id}$. Restricting to the flat irreducible connections gives the fibration

$$(4.1) \quad \mathcal{G}/\pm \text{Id} \rightarrow \mathcal{F}^* \rightarrow \mathcal{M}^*.$$

That $\mathcal{M}^*$ is simply connected follows immediately from [13, Theorem 7.1] and the fact that $\mathcal{M}^*$ is homeomorphic to the space of stable rank 2 holomorphic bundles over Σ with trivial determinant (see [24, 1]).

The calculations in [13] show that $\pi_2(\mathcal{M}^*) = \mathbb{Z} \oplus \mathbb{Z}/2$ and it is not hard to show that $\pi_1(\mathcal{G}/\pm \text{Id}) = \mathbb{Z} \oplus \mathbb{Z}/2$. If we knew that the map induced by the connecting homomorphism in the long exact sequence of homotopy groups for the fibration (4.1) were an isomorphism (or merely surjective), then the result would follow from this long exact sequence since $\pi_1(\mathcal{M}^*) = 0$. This is true, in fact the results of [13], though stated for $U(2)$ instead of $SU(2)$, are equally valid for $SU(2)$.

Alternatively, we can show that $\pi_1(\mathcal{F}^*)$ is zero using the result of [26]. Fix a flat irreducible connection $A \in \mathcal{F}^*$. Let $\gamma : S^1 \rightarrow \mathcal{F}$ be a loop of flat connections. Corollary 2.7 of [14] shows that $\mathcal{A}^s$ is simply connected, and since $\mathcal{F}^* \subset \mathcal{A}^s$ (see [1]) γ bounds a disc $\bar{\gamma} : D^2 \rightarrow \mathcal{A}^s$. The main result of [26] asserts that the Yang-Mills flow on $\mathcal{A}$, restricted to $\mathcal{A}^s$, defines a strong deformation retraction of $\mathcal{A}^s$ to $\mathcal{F}^*$. Thus γ bounds a disc in $\mathcal{F}^*$, completing the proof. $\square$

The space of irreducible $SU(2)$ representations of a free group is easily proven to be path connected. Thus the space $\mathcal{M}^*(X_1)$, homeomorphic to the space of conjugacy classes of irreducible representations of $\pi_1(X_1)$, is path connected. There is a fibration

$$\mathcal{G}(X_1)/\pm \text{Id} \rightarrow \mathcal{F}^*(X_1) \rightarrow \mathcal{M}^*(X_1)$$

as in the proof of Lemma 4.1. The gauge group $\mathcal{G}(X_1) = \text{Map}(X_1, SU(2))$ is path connected since X_1 has the homotopy type of a 1-complex. Thus $\mathcal{F}^*(X_1)$ is path connected. The same argument shows that the space of irreducible perturbed flat connections on X_2 is also path connected, since the perturbation was chosen small enough so that $\mathcal{M}_f^*(X_2)$ is diffeomorphic to the unperturbed moduli space $\mathcal{M}^*(X_2)$.

Combining the facts of the preceding paragraph, Lemma 4.1, and the observation that since the space of connections on a manifold is contractible, any family of connections on a submanifold of X extends to a family of connections on all of X , we obtain the following useful proposition.

Proposition 4.2. *Given perturbed flat irreducible $SU(2)$ connections A_0, A_1 on the closed 3-manifold X , there exists a 2-parameter family $A_{s,t}$, $s, t \in [0, 1]$ of irreducible connections on X satisfying:*

- (1) $A_{0,t} = A_0$ and $A_{1,t} = A_1$ for all $t \in [0, 1]$.
- (2) $A_{s,0}$ restricts to a path of irreducible flat connections on X_1 .
- (3) $A_{s,1}$ restricts to a path of irreducible perturbed flat connections on X_2 .
- (4) $A_{s,t}$ restricts to an irreducible flat connection $a_{s,t}$ on Σ for all s, t .
- (5) The restriction of $A_{s,t}$ to a collar of Σ is in product form: $A_{s,t}|_{\Sigma \times [-1,1]} = \frac{\partial}{\partial u} + a_{s,t}$. $\square$

We now give an argument that $\text{Sign}_T = \text{Sign}_C$. The strategy is to identify Sign_T with an infinite-dimensional Maslov index of the Cauchy data spaces, then to use homotopy and stretching arguments to symplectically reduce to a finite-dimensional Maslov index which we identify with Sign_C . We

recommend that the reader who is unfamiliar with the type of argument we use look at the “user’s guide” section of [12].

Let $\Lambda_{s,t}^1$ denote the Cauchy data space for the restriction of the operator $D_{A_{s,t},f}$ to X_1 . Similarly define $\Lambda_{s,t}^2$. These vary continuously by Theorem 3.19. Note that in contrast to Section 3, the superscripts 1 and 2 refer to the Cauchy data spaces for the two parts of the Heegard decomposition, not to the Sobolev indices. Thus $\Lambda_{s,t}^1$ and $\Lambda_{s,t}^2$ are continuous families of Lagrangian subspaces of $L^2(\Omega_\Sigma^*(\text{ad } P))$. Moreover, for each (s,t) , $(\Lambda_{s,t}^1, \Lambda_{s,t}^2)$ is a Fredholm pair. Hence the Maslov index $\text{Mas}(\Lambda_\alpha^1, \Lambda_\alpha^2) \in \mathbb{Z}$ is defined for any path $\alpha : [0, 1] \rightarrow [0, 1] \times [0, 1]$ (for details see [22]).

Nicolaescu’s theorem says that the spectral flow of the family $D_{A_{s,0},f}$, $s \in [0, 1]$ equals the (infinite-dimensional) Maslov index $\text{Mas}(\Lambda_{s,0}^1, \Lambda_{s,0}^2)$ (see [25, 22]). Hence

$$\text{Sign}_T([A_0], [A_1]) = (-1)^{\text{SF}(D_{A_{s,0}})} = (-1)^{\text{Mas}(\Lambda_{s,0}^1, \Lambda_{s,0}^2)}.$$

We will use the homotopy properties of the Maslov index to find a simpler, finite-dimensional expression for $\text{Mas}(\Lambda_{s,0}^1, \Lambda_{s,0}^2)$ which can be compared to the difference in intersection numbers.

The reduction will be based on the following observations. Since the connection $a_{s,t}$ on Σ is flat for all s, t , the kernel of $S_{a_{s,t}}$ is identified via the Hodge theorem with the cohomology $H^*(\Sigma; \text{ad } P_{a_{s,t}})$ of the twisted de Rham complex

$$(4.2) \quad 0 \rightarrow \Omega^0(\Sigma, \text{ad } P_{a_{s,t}}) \xrightarrow{d_{a_{s,t}}} \Omega^1(\Sigma, \text{ad } P_{a_{s,t}}) \xrightarrow{d_{a_{s,t}}} \Omega^2(\Sigma, \text{ad } P_{a_{s,t}}) \rightarrow 0.$$

Since $a_{s,t}$ is irreducible, $H^0(\Sigma; \text{ad } P_{a_{s,t}}) = 0$ (i.e. using (2.3) for Σ) and so $H^2(\Sigma; \text{ad } P_{a_{s,t}}) = 0$ by Poincaré duality. Thus the kernel of $S_{a_{s,t}}$ is just $H^1(\Sigma; \text{ad } P_{a_{s,t}})$. But this is canonically identified with the tangent space of $\mathcal{M}^*(\Sigma)$ at $[a_{s,t}]$, since the image of $d_{a_{s,t}} : \Omega^0(\Sigma, \text{ad } P_{a_{s,t}}) \rightarrow \Omega^1(\Sigma, \text{ad } P_{a_{s,t}})$ is the tangent space of the $\mathcal{G}(\Sigma)$ orbit and $d_{a_{s,t}} : \Omega^1(\Sigma, \text{ad } P_{a_{s,t}}) \rightarrow \Omega^2(\Sigma, \text{ad } P_{a_{s,t}})$ is the linearization at $a_{s,t}$ of the curvature map $a \mapsto F(a)$.

Via this identification, the cup product (or wedge product if one identifies cohomology with harmonic forms)

$$H^1(\Sigma; \text{ad } P_{a_{s,t}}) \times H^1(\Sigma; \text{ad } P_{a_{s,t}}) \xrightarrow{\cup} H^2(\Sigma; \mathbb{R}) = \mathbb{R}$$

defines a symplectic structure on $\mathcal{M}^*(\Sigma)$. In particular, since $\mathcal{M}^*(\Sigma)$ is a smooth $6g - 6$ dimensional manifold, $\ker S_{a_{s,t}} = H^1(\Sigma; \text{ad } P_{a_{s,t}})$ is a smoothly varying family of symplectic subspaces of $L^2(\Omega_\Sigma^*(\text{ad } P))$. For convenience we use the notation

$$H_{s,t}^1(\Sigma) = H^1(\Sigma; \text{ad } P_{a_{s,t}}).$$

Let $P_{s,t}^+$ denote the (L^2 closure of the) positive eigenspan of the tangential operator $S_{a_{s,t}}$, and let $P_{s,t}^-$ denote its negative eigenspan. Thus we have a smoothly varying (in s, t) decomposition

$$L^2(\Omega_\Sigma^*(\text{ad } P)) = P_{s,t}^- \oplus H_{s,t}^1(\Sigma) \oplus P_{s,t}^+.$$

Lemma 4.3. *For each $s \in [0, 1]$, let $T_{s,0}^1 \subset H_{s,0}^1(\Sigma)$ be the subspace defined by*

$$T_{s,0}^1 = \text{image } H^1(X_1; \text{ad } P_{a_{s,0}}) \rightarrow H^1(\Sigma; \text{ad } P_{a_{s,0}}).$$

Then the spaces $T_{s,0}^1$ are Lagrangian and vary continuously. Via the identification $T_{a_{s,0}} \mathcal{M}^(\Sigma) \cong H_{s,0}^1(\Sigma)$ the subspace $T_{s,0}^1$ corresponds to the tangent space at $a_{s,0}$ of the submanifold $\mathcal{M}^*(X_1)$ of $\mathcal{M}^*(\Sigma)$.*

Moreover

$$\text{SF}(D_{A_{s,0},f}) = \text{Mas}(\Lambda_{s,0}^1, \Lambda_{s,0}^2) = \text{Mas}(P_{s,0}^+ \oplus T_{s,0}^1, \Lambda_{s,0}^2).$$

Proof. The statement of Lemma 4.3 refers only to the parameters $(s, 0)$, $s \in [0, 1]$; in other words, throughout the proof the “ t ” parameter is zero.

The fact that $T_{s,0}^1$ is Lagrangian is a standard fact in geometric topology whose easy proof (based on Poincaré duality) we leave to the reader. The identification of $T_{s,0}^1$ with the tangent space of $\mathcal{M}^*(X_1)$ is also standard; one of many proofs is to consider the map of de Rham complexes

$\Omega^*(X_1; \text{ad } P) \rightarrow \Omega^*(\Sigma; \text{ad } P)$ in light of the diagram (2.3). Since $\mathcal{M}^*(X_1)$ is a smooth submanifold, it follows that $T_{s,0}^1$ varies smoothly. Thus we focus on the statements about the Maslov index.

It follows from the adiabatic limit theorem of [25] and its refinement in [12] that replacing the collar of X_1 by an increasingly longer collar $\Sigma \times [-R, 0]$ gives a homotopy of the path $s \mapsto \Lambda_{s,0}^1$ to $s \mapsto P_{s,0}^+ \oplus T_{s,0}^1$. This is a consequence of the fact that there are no L^2 solutions to $D_{A_{s,0},f}\phi = 0$ on $(\Sigma \times (-\infty, 0]) \cup X_1$, which follows from [2] and two facts:

- (1) The restriction map $H^1(X_1; \text{ad } P_{a_{s,0}}) \rightarrow H^1(\Sigma; \text{ad } P_{a_{s,0}})$ is injective.
- (2) The space $T_{s,0}^1$ equals the *limiting values of extended L^2 solutions* in the sense of [2]. This was first observed by Yoshida [31] and in the present notation a proof can be found in [12] and [22].

Let $r \mapsto \Lambda_{s,0}^1(r)$ denote this homotopy. To be explicit,

$$\Lambda_{s,0}^1(r) = \begin{cases} \text{the Cauchy data space for } D_{A_{s,0},f} \text{ on } (\Sigma \times [-\frac{1}{1-r}, 0]) \cup X_1 & \text{if } r < 1, \\ P_{s,0}^+ \oplus T_{s,0}^1 & \text{if } r = 1. \end{cases}$$

This is not a homotopy rel endpoints, and so to see why the Maslov indices nevertheless agree, it suffices to show that the dimensions of the intersections $\Lambda_{0,0}^1(r) \cap \Lambda_{0,0}^2$ and $\Lambda_{1,0}^1(r) \cap \Lambda_{1,0}^2$ are independent of r , in fact these intersections are zero for all r .

For $r < 1$ these intersections are isomorphic to the kernel of the extension of $D_{A_{i,0},f}$ ($i = 0, 1$) to $X(r) = (\Sigma \times [-\frac{1}{1-r}, 0]) \cup X_1 \cup X_2$, i.e. X with a long collar inserted near Σ . But the dimension of the kernel of $D_{A_{i,0},f}$ is independent of the choice of Riemannian metric on X since $A_{i,0}$ is perturbed flat: it is isomorphic to the cohomology of the complex (2.7), which is zero by the assumption that each perturbed flat connection is non-degenerate.

For $r = 1$, we need to see that $(P_{0,0}^+ \oplus T_{0,0}^1) \cap \Lambda_{0,0}^2$ and $(P_{1,0}^+ \oplus T_{1,0}^1) \cap \Lambda_{1,0}^2$ are both zero. The quotient

$$\frac{(P_{0,0}^+ \oplus \ker S_{a_{0,0}}) \cap \Lambda_{0,0}^2}{P_{0,0}^+ \cap \Lambda_{0,0}^2} \subset \ker S_{a_{0,0}}$$

is just $T_{0,0}^2$ ([2, 22]) and the fact that each perturbed flat connection is non-degenerate implies that $T_{0,0}^1 \cap T_{0,0}^2 = 0$. Moreover, $P_{0,0}^+ \cap \Lambda_{0,0}^2 = 0$ since any section in this intersection is the restriction to the boundary of a solution to $D_{A_{0,0},f}\phi = 0$ which extends to an L^2 solution on $(\Sigma \times (-\infty, 0]) \cup X_1$. The only such ϕ is zero, as we argued above. Hence $(P_{0,0}^+ \oplus T_{0,0}^1) \cap \Lambda_{0,0}^2 = 0$. The other endpoint is handled identically.

Since the Maslov index is unchanged when the paths of Lagrangians are homotoped by a homotopy that preserves the dimension of the intersections at the endpoints ([22]),

$$\text{Mas}(\Lambda_{s,0}^1, \Lambda_{s,0}^2) = \text{Mas}(P_{s,0}^+ \oplus T_{s,0}^1, \Lambda_{s,0}^2),$$

as desired. $\square$

The spaces $H_{s,t}^1 = \ker S_{s,t}$ form a symplectic vector bundle over the square $[0, 1] \times [0, 1]$. In fact, this is just the pullback of the tangent bundle $T_*\mathcal{M}^*(\Sigma)$ via the map $[0, 1] \times [0, 1] \rightarrow \mathcal{M}^*(\Sigma)$ given by $(s, t) \mapsto [a_{s,t}]$.

We have a Lagrangian subbundle $T_{s,0}^1$ defined over the top edge of this square, and also the 2 sides since $A_{0,t}$ is independent of t and $A_{1,t}$ is independent of t . Extend this arbitrarily (e.g. pull back via a retraction of the square to its three sides) to a Lagrangian subbundle $T_{s,t}^1$ over the entire square. Then the 2-parameter family $P_{s,t}^+ \oplus T_{s,t}^1$ provides a homotopy which shows that

$$(4.3) \quad \text{Mas}(P_{s,0}^+ \oplus T_{s,0}^1, \Lambda_{s,0}^2) = \text{Mas}(P_{s,1}^+ \oplus T_{s,1}^1, \Lambda_{s,1}^2).$$

Together with Lemma 4.3, (4.3) implies that

$$(4.4) \quad SF(D_{A_{s,0},f}) = \text{Mas}(P_{s,1}^+ \oplus T_{s,1}^1, \Lambda_{s,1}^2).$$

An argument very similar to the proof of Lemma 4.3, this time taking the adiabatic limit on X_2 , shows that

$$(4.5) \quad \text{Mas}(P_{s,1}^+ \oplus T_{s,1}^1, \Lambda_{s,1}^2) = \text{Mas}(P_{s,1}^+ \oplus T_{s,1}^1, P_{s,1}^- \oplus T_{s,1}^2),$$

using the fact that the path of connections $A_{s,1}$ is perturbed flat on X_2 . Moreover, $T_{s,1}^2 \subset H_{s,1}^1(\Sigma)$ is isomorphic to image of the differential of the inclusion $\mathcal{M}_f^*(X_2) \rightarrow \mathcal{M}^*(\Sigma)$ at $A_{s,1}$, i.e. $T_{s,1}^2 = \text{image}(T_{A_{s,1}}(\mathcal{M}_f^*(X_2)) \rightarrow T_{A_{s,1}}(\mathcal{M}^*(\Sigma)))$. Combining (4.4) and (4.5) and the facts that the Maslov index is additive with respect to direct sum of symplectic spaces and that $P_{s,1}^+$ and $P_{s,1}^-$ are always orthogonal, one arrives at the conclusion

$$(4.6) \quad \text{SF}(D_{A_{s,0},f}) = \text{Mas}(T_{s,1}^1, T_{s,1}^2).$$


The spectral flow $\text{SF}(D_{A_{s,0},f})$ depends only on f and the endpoints A_0 and A_1 of A_t ; see Lemma 2.1. Moreover its reduction modulo 2 (in fact modulo 8) only depends on the gauge equivalence classes $[A_0]$ and $[A_1]$. Equation (4.6) has identified $\text{SF}(D_{A_{s,0},f})$ with a finite-dimensional Maslov index corresponding to the tangent spaces of Lagrangian submanifolds of a symplectic manifold. The following lemma identifies the mod 2 reduction of this Maslov index with the difference in intersection numbers of $\mathcal{M}^*(X_1)$ and $\mathcal{M}^*(X_2)$ at $[A_0]$ and $[A_1]$.

Lemma 4.4. *Let A and B be smooth oriented Lagrangian submanifolds of a (finite-dimensional) symplectic manifold W . Let p and q be two transverse intersection points of A and B . Let γ_A be a path in A from p to q and γ_B a path in B from p to q . Suppose that $F(s,t)$ is a homotopy rel endpoints in W from γ_A to γ_B . Let $T_{s,0}^A$ be the Lagrangian subbundle of the restriction of T_*W to γ_A consisting of tangent vectors to A , and similarly let $T_{s,1}^B$ be the Lagrangian subbundle of the restriction of T_*W to γ_B consisting of tangent vectors to B .*

Then T^A extends over the homotopy, and if $T_{s,t}^A$ is any extension (with $T_{0,s}^A$ and $T_{1,t}^A$ independent of t), then $\text{Mas}(T_{s,1}^A, T_{s,1}^B)$ is even if and only if intersection number of A and B at p and q coincide.

□

The straightforward proof can easily be constructed by musing on the following picture, and is left to the reader.



Lemma 4.4 and Equation (4.6) then imply Taubes' theorem.

Theorem 4.5 (Taubes). *$\text{Sign}_C = \text{Sign}_T$, and hence Casson's and Taubes' invariants are equal up to an overall sign (which may depend on the homology 3-sphere).*

□

Taubes proves more, namely he gives a construction for specifying an overall sign so that in fact his invariant equals Casson's invariant for all homology 3-spheres.

REFERENCES

- [1] ATIYAH, M. F. and BOTT, R. *The Yang-Mills equations over Riemann surfaces*. Philos. Trans. Roy. Soc. London Ser. A 308 (1983), no. 1505, 523–615.
- [2] ATIYAH, M. F., PATODI, V. K. and SINGER, I. M. *Spectral asymmetry and Riemannian geometry. I*. Math. Proc. Cambridge Philos. Soc. 77 (1975), 43–69.
- [3] BODEN, HANS U. and HERALD, CHRISTOPHER M. *The $SU(3)$ Casson invariant for integral homology 3-spheres*. J. Differential Geom. 50 (1998), no. 1, 147–206.
- [4] BODEN, HANS U., HERALD, CHRISTOPHER M., and KIRK, PAUL *An integer valued $SU(3)$ Casson invariant*. Math. Res. Lett. 8 (2001), no. 5-6, 589–603.
- [5] BODEN, HANS U., HERALD, CHRISTOPHER M., KIRK, PAUL, and KLASSEN, ERIC P. *Gauge theoretic invariants of Dehn surgeries on knots*. Geom. Topol. 5 (2001), 143–226.
- [6] BODEN, HANS U., HERALD, CHRISTOPHER M., and KIRK, PAUL *The $SU(3)$ Casson Invariant for Brieskorn spheres*. In preparation.
- [7] BOOSS-BAVNBEEK, B. *The unique continuation property for Dirac operators revisited*. Geometry and topology: Aarhus (1998), Contemp. Math. 258 (2000), 21–32.
- [8] BOOSS-BAVNBEEK, B., LESCH, MATTHIAS, and PHILLIPS, JOHN *Unbounded Fredholm operators and spectral flow*. Preprint, math.FA/0108014.
- [9] BOOSS-BAVNBEEK, B., MARCOLLI, M., and WANG, B.L. *Weak UCP and perturbed monopole equations*. Preprint, math.DG/0203171.
- [10] BOOSS-BAVNBEEK, BERNHELM, and WOJCIECHOWSKI, KRZYSZTOF P. *Elliptic boundary problems for Dirac operators*. Mathematics: Theory & Applications. Birkhäuser Boston, Inc., Boston, MA, 1993. xviii+307 pp.
- [11] CAPPELL, S., LEE, R., and MILLER, E. Y. *A perturbative $SU(3)$ Casson invariant*. Comment. Math. Helv. 77 (2002), no. 3, 491–523.
- [12] DANIEL, A. M. and KIRK, PAUL *A general splitting formula for the spectral flow*. With an appendix by K. P. WOJCIECHOWSKI. Michigan Math. J. 46 (1999), no. 3, 589–617.
- [13] DASKALOPOULOS, GEORGIOS D. *The topology of the space of stable bundles on a compact Riemann surface*. J. Differential Geom. 36 (1992), no. 3, 699–746.
- [14] DASKALOPOULOS, GEORGIOS D. and UHLENBECK, KAREN K. *An application of transversality to the topology of the moduli space of stable bundles*. Topology 34 (1995), no. 1, 203–215.
- [15] DONALDSON, S. K. and KRONHEIMER, P. B. *The geometry of four-manifolds*. Oxford Mathematical Monographs. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1990. x+440 pp.
- [16] DOSTOGLU, STAMATIS and SALAMON, DIETMAR A. *Self-dual instantons and holomorphic curves*. Ann. of Math. (2) 139 (1994), no. 3, 581–640.
- [17] FLOER, ANDREAS *An instanton-invariant for 3-manifolds*. Comm. Math. Phys. 118 (1988), no. 2, 215–240.
- [18] HERALD, CHRISTOPHER M. *Legendrian cobordism and Chern-Simons theory on 3-manifolds with boundary*. Comm. Anal. Geom. 2 (1994), no. 3, 337–413.
- [19] HERALD, CHRISTOPHER M. in preparation.
- [20] FREED, DANIEL S. and UHLENBECK, KAREN K. *Instantons and four-manifolds*. Second edition. Mathematical Sciences Research Institute Publications, 1. Springer-Verlag, New York, 1991. xxii+194 pp.
- [21] GRUBB, GERD *Trace expansions for pseudodifferential boundary problems for Dirac-type operators and more general systems*. Ark. Math. 37 (1999), 45–86.
- [22] KIRK, PAUL and LESCH, MATTHIAS *The eta-invariant, Maslov index, and spectral flow for Dirac-type operators on manifolds with boundary*. Preprint, math.DG/0012123.
- [23] LIONS, J. and MAGENES, E. *Non-homogeneous boundary problems and applications*. Springer-Verlag, New York, 1972.
- [24] NARASIMHAN, M. S. and SESHADRI, C. S. *Stable and unitary vector bundles on a compact Riemann surface*. Ann. of Math. (2) 82 1965 540–567.
- [25] NICOLAESCU, LIVIU I. *The Maslov index, the spectral flow, and decompositions of manifolds*. Duke Math. J. 80 (1995), no. 2, 485–533.
- [26] RADE, JOHAN *On the Yang-Mills heat equation in two and three dimensions*. J. Reine Angew. Math. 431 (1992), 123–163.
- [27] RAMADAS, T. R., SINGER, I. M., and WEITSMAN, J. *Some comments on Chern-Simons gauge theory*. Comm. Math. Phys. 126 (1989), no. 2, 409–420.
- [28] SCOTT, S. G. *Determinants of Dirac boundary value problems over odd-dimensional manifolds*. Commun. Math. Phys. 173 (1995), 43–76.
- [29] TAUBES, CLIFFORD HENRY *Casson's invariant and gauge theory*. J. Differential Geom. 31 (1990), no. 2, 547–599.
- [30] TAYLOR, MICHAEL E. *Partial differential equations*. Applied Mathematical Sciences Vol 115, Springer-Verlag, New York, 1996.
- [31] YOSHIDA, TOMOYOSHI *Floer homology and splittings of manifolds*. Ann. of Math. (2) 134 (1991), no. 2, 277–323.

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, BLOOMINGTON, IN 47405
E-mail address: bhimpel@indiana.edu
URL: <http://mypage.iu.edu/~bhimpel>

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, BLOOMINGTON, IN 47405
E-mail address: pkirk@indiana.edu
URL: <http://mypage.iu.edu/~pkirk>

UNIVERSITÄT ZU KÖLN, MATHEMATISCHES INSTITUT, WEYERTAL 86–90, D–50931 KÖLN
E-mail address: lesch@mi.uni-koeln.de
URL: <http://www.mi.uni-koeln.de/~lesch>